

PRIMARY TYPE SPECIMENS OF MARINE MOLLUSCA  
(EXCLUDING CEPHALOPODA)  
IN THE SOUTH AFRICAN MUSEUM

By

ELIZABETH GILES

&

TERRENCE GOSLINER

*South African Museum, Cape Town*

[MS accepted 2 November 1982]

ABSTRACT

A list of type specimens of marine Mollusca contained in the collections of the South African Museum is presented. Three hundred and thirty-seven species are included in this list.

CONTENTS

	PAGE
Introduction . . . . .	1
Polyplacophora . . . . .	2
Gastropoda . . . . .	3
Scaphopoda . . . . .	40
Bivalvia . . . . .	41
References . . . . .	49

INTRODUCTION

The molluscan collections housed in the South African Museum have provided the basis for numerous systematic investigations and extensive studies by well-known malacologists such as G. B. Sowerby, J. R. Tomlin, and W. H. Turton. In 1958 K. H. Barnard initiated his work on the molluscan fauna, based largely on material obtained from the dredging operations of the R.S. *Pieter Faure*.

As a result of these research activities spanning more than a century, many primary type specimens have been deposited in the collections of the South African Museum. In his posthumously published faunal list of South African molluscs Barnard (1974) noted some of the type material but a complete list has not been published.

The designation of type specimens was not indicated in most of the works of the above-mentioned authors and their status has been largely determined on the basis of review of K. H. Barnard's research notes, which involve correspondence with most of his predecessors as well as his own work.

Localities and data are given as they appear in the original publications or have been supplemented by information contained in catalogues of the molluscan holdings in the South African Museum.

## SPECIES LIST

## Class POLYPLACOPHORA

Family *Acanthochitonidae*

- Acanthochiton turtoni* Ashby, 1928: 79, pl. 6 (figs 1–4)  
= *Acanthochitona garnoti* (Blainville): Barnard, 1963*d*: 328.  
Holotype: SAM–A32649. Valve of holotype.  
Locality: Port Alfred; no depth or date.  
Collected by: Col. W. Turton.

Family *Lepidopleuridae*

- Lepidopleurus chariessa* Barnard, 1963*d*: 332, fig. 29h.  
Syntypes: SAM–A9338. Two shells, disarticulated.  
Locality: off Cape St. Blaize; 125 fm.; 1899.  
Collected by: R.S. Pieter Faure.

Family *Chitonidae*

- Chiton ashbyi* Barnard, 1963*d*: 343, fig. 29m.  
Holotype: SAM–A9337. Middle valve of holotype.  
Locality: off Cape Infanta; 46 fm.; 10 July 1900.  
Collected by: R.S. Pieter Faure.
- Chiton barnardi* Ashby, 1931: 46, pl. 6 (figs 74–76), pl. 7 (fig. 77).  
Holotype: SAM–A5331. Shell.  
Paratype: SAM–A6590. One valve.  
Locality: Mozambique Island (?Inhaca); no depth; 1912.  
Collected by: K. H. Barnard.
- Chiton (Hanleya) sykesi* Sowerby, 1903: 225, pl. 5 (fig. 13)  
= *Lepidopleurus sykesi* (Sowerby): Ashby, 1931: 15, pl. 2 (figs 16–19).  
Syntypes: SAM–5342. Two, girdle on slide.  
Locality: off Cape Point; 210 fm.; 11 June 1900.  
Collected by: R.S. Pieter Faure.

Family *Ischnochitonidae*

- Chaetopleura pomarium* Barnard, 1963*d*: 335, fig. 29i.  
Holotype: SAM–A9339. Shell with animal.  
Locality: western Cape Peninsula; no depth or date.  
Collected by: University of Cape Town.

*Ischnochiton delagoaensis* Ashby, 1931: 40, pl. 6 (figs 63–66).

Holotype: SAM–A6589. Valves, disarticulated.

Locality: Delagoa Bay; no depth; 1912.

Collected by: K. H. Barnard.

*Ischnochiton hewitti* Ashby, 1931: 33, pl. 5 (figs 50–53).

Holotype: SAM–A6757. Median valve of holotype.

Paratypes: SAM–A33953. Eight valves.

Locality: Mouille Point, Table Bay; intertidal; 1900.

Collected by: R. M. Lightfoot.

#### Class GASTROPODA

#### Subclass PROSOBRANCHIA

#### Family Pleurotomariidae

*Pleurotomaria africana* Tomlin, 1948: 2, pl. 1.

Holotype: SAM–A29663. Shell.

Locality: 29°48'S 31°23'E; 366 m; 21 July 1931.

Collected by: R.S. *Africana II*.

#### Family Scissurellidae

*Scissurella africanae* Barnard, 1963*b*: 13.

Holotype: SAM–A29933. Shell.

Locality: 30°48'S 52°08'E; 400 m; 9 July 1961.

Collected by: R.S. *Africana II*.

*Scissurella tabulata* Barnard, 1964*c*: 21.

Holotype: SAM–A9335 (not A9345). Shell with operculum.

Locality: off Cape Natal; 804 m; 4 April 1901.

Collected by: R.S. *Pieter Faure*.

#### Family Fissurellidae

*Glyphis spreta* Smith, 1901: 104, pl. 1 (fig. 18)

= *Diodora spreta* (Smith): Tomlin, 1932: 160.

Syntypes: SAM–2459. Three shells.

Locality: Mossel Bay; intertidal; 1898.

Collected by: Miss Powrie.

*Emarginula koon* Kilburn, 1978: 439, pl. 5.

Paratype: SAM–A33924. Shell.

Locality: Shelly Beach near Port Shepstone; November 1965.

Collected by: R. Cock.

*Emarginula natalensis* Barnard, 1963d: 297, fig. 23.

Syntypes: SAM-A9325. Three shells, worn.

Locality: off Cape Morgan; 141 m; 26 July 1901.

Collected by: R.S. Pieter Faure.

*Emarginula oppressa* Barnard, 1963d: 298, fig. 24a

= *Emarginella oppressa* (Barnard): Kilburn, 1978: 446, pl. 8 (fig. b).

Holotype: SAM-A9330. Shell.

Locality: off Cape Natal; 98 m; 14 December 1900.

Collected by: R.S. Pieter Faure.

*Emarginula pulchreclathrata* Tomlin, 1932: 162, fig. 4

= *Emarginula agulhasensis* Thiele: Kilburn, 1978: 433, pl. 2a-b.

Holotype: SAM-A3617. Shell.

Locality: Saldanha Bay; 98 m; 13 March 1902.

Collected by: R.S. Pieter Faure.

*Emarginula thorektes* Kilburn, 1978: 440, pl. 6a-c.

Paratype: SAM-A9323. Shell.

Locality: off Cape Morgan; 141 m; 26 July 1901.

Collected by: R.S. Pieter Faure.

*Emarginula vadum* Barnard, 1963d: 297, figs 23d-24b.

Holotype: SAM-A9329. Shell.

Locality: off Cape Vidal; 146-183 m; 14 December 1900.

Collected by: R.S. Pieter Faure.

*Parmaphorella barnardi* Tomlin, 1932: 164, fig. 5

= *Tugali barnardi* (Tomlin): Kilburn, 1978: 452, pl. 11 (figs b-d).

Holotype: SAM-A3623. Shell.

Locality: off Cape Point; 344 m; 27 February 1902.

Collected by: R.S. Pieter Faure.

*Puncturella (Vacerrrena) christiaensi* Kilburn, 1978: 448, pl. 9 (figs c-e).

Paratype: SAM-A33925. Shell.

Locality: off Sodwana Bay; 50 m; 1975.

Collected by: R.V. Meiring Naude.

#### Family Trochidae

*Angaria lacunosa* Barnard, 1963d: 283.

Holotype: SAM-A9281. Shell.

Locality: off Natal; no depth or date.

Collected by: R.S. Pieter Faure.

*Bassilissa gelida* Barnard, 1963b: 440, fig. 8b.

Holotype: SAM-A9720. Shell, radula, slide.

Locality: 33°26'S 16°33'E; 2 380 m; August 1959.

Collected by: F. H. Talbot on board R.S. *Africana II*.

*Calliostoma circus* Barnard, 1969: 653, fig. 27b.

Holotype: SAM-A30032. Shell with operculum, radula slide.

Locality: off Cape Point; no depth or date.

Collected by: commercial trawler.

*Calliostoma glaucophaos* Barnard; 1963b: 442, fig. 8c.

Syntypes: SAM-A9830. Two shells with opercula, radula slide.

Locality: 33°52'S 16°51'E; 2 520–2 810 m; December 1959.

Collected by: F. H. Talbot on board R.S. *Africana II*.

*Calliostoma (Lischkeia) granoliratum* Sowerby, 1903: 222, pl. 5 (fig. 7)

= *Calliotropis granolirata* (Sowerby): Thiele, 1925: 48.

Holotype: SAM-A5296. Shell with operculum.

Paratype: SAM-A33690 (previously A5296). Shell.

Locality: off Cape Point; 665 m; 12 June 1900.

Collected by: R.S. *Pieter Faure*.

*Calliostoma iridescens* Sowerby, 1903: 223, pl. 5 (fig. 4).

Holotype: SAM-A5294. Shell.

Locality: off Cape Natal; 101 m; 24 December 1900.

Collected by: R.S. *Pieter Faure*.

*Calliostoma perfragile* Sowerby, 1903: 222, pl. 5 (fig. 3).

Holotype: SAM-A5275. Shell.

Locality: off Cape Point; 282–304 m; 18 April 1900.

Collected by: R.S. *Pieter Faure*.

*Calliotropis chenoderma* Barnard, 1963d: 263, fig. 16c.

Syntype: SAM-A9296. Shell.

Locality: off Cape St. Blaize; 229 m; 21 December 1899.

Syntype: SAM-A9297. Shell.

Locality: off Cape Morgan; 86 m; 25 July 1901.

Collected by: R.S. *Pieter Faure*.

*Calliotropis persculpta* (Sowerby): see *Solariella persculpta* p. 7.

*Calliotropis pompe* Barnard, 1963c: 440, fig. 8b.

Syntypes: SAM-A9883. Two shells, one with operculum.

Locality: off Cape Point, 34°42'S 16°54'E; 1 725–1 789 fm.; December 1959.

Syntype: SAM-A9795. Shell.

Locality: off Cape Point, 33°50'S 16°30'E; 1 450–1 660 fm.; December 1959.

Collected by: F. H. Talbot on board R.S. *Africana II*.

*Gibbula beckeri* Sowerby, 1901: 214, pl. 22 (fig. 22).

Paratype: SAM-A3321. Shell.

Locality: Port Alfred; intertidal; 1915.

Collected by: H. Becker.

*Gibbula (Enida) perspectiva* Sowerby, 1900: 6, pl. 1 (fig. 17).

Paratype: SAM-A3320. Shell.

Locality: Pondoland; intertidal; 1915.

Collected by: H. Becker.

*Guttula blanda* Barnard, 1963d: 265, fig. 17.

Holotype: SAM-A7424. Shell, radula slide.

Locality: off Cape Point; 1 270 m; 20 August 1903.

Collected by: R.S. *Pieter Faure*.

*Minolia adarticulata* Barnard, 1963d: 235, fig. 11f.

Syntypes: SAM-A9277. Two shells.

Locality: off Hood Point, East London; 90 m; 15 July 1901.

Collected by: R.S. *Pieter Faure*.

*Minolia cycloma* Barnard, 1964c: 18, fig. 2c–d.

Holotype: SAM-A9279. Shell.

Locality: off Tugela River, Natal; 67 m; 9 January 1901.

Collected by: R.S. *Pieter Faure*.

*Minolia congener* Sowerby, 1903: 223, pl. 5 (fig. 2)

= *Solariella laevis* (von Martens): Barnard, 1963d: 239, fig. 10c.

Syntypes: SAM-A5257. Two shells with opercula.

Locality: off Cape St. Blaize; 67 m; 22 October 1900.

Collected by: R.S. *Pieter Faure*.

*Seguenzia fatigans* Barnard, 1963b: 13, pl. 2 (figs 10–11).

Syntypes: SAM-A29930. Two shells.

Locality: 27°48'S 47°19'E; 875 m; 22 June 1961.

Collected by: R.S. *Africana II*.

*Seguenzia simplex* Barnard, 1963d: 265, fig. 16a.

Syntypes: SAM-A7421. Four shells (one juvenile); radula slide.

Locality: off Cape Point; 1 280 m; 20 August 1903.

Collected by: R.S. *Pieter Faure*.

- Solariella beckeri* Sowerby, 1901: 214, pl. 22 (fig. 21)  
= *Solariella laevissima* (von Martens): Barnard, 1963d: 239, fig. 10c.  
Paratype: SAM-A3323. Shell.  
Locality: Kowie (Port Alfred); no depth; 1915.  
Collected by: H. Becker.
- Solariella dowi* Barnard, 1963b: 13, pl. 2 (figs 10–11).  
Syntypes: SAM-A29932. Two shells.  
Locality: 27°48'S 47°19'E; 875 m; 22 June 1961.  
Collected by: R.S. *Africana II*.
- Solariella franciscana* Barnard, 1963d: 243, figs 10e–11b  
= *Spectamen franciscana* (Barnard): Kilburn, 1977: 178.  
Holotype: SAM-A3615. Shell with operculum, radula slide.  
Locality: off Cape St. Francis; 137 m; 19 February 1902.  
Collected by: R.S. *Pieter Faure*.
- Solariella gilchristi* Barnard, 1963d: 242, fig. 11d.  
Syntypes: SAM-A3605. Two shells.  
Locality: off Buffalo River; 564 m; 24 April 1901.  
Collected by: R.S. *Pieter Faure*.
- Solariella macleari* Barnard, 1963d: 241, fig. 11c.  
Syntypes: SAM-A7417. Four shells, one apex, two opercula, radula slide.  
Locality: 'off Cape Point N 89°E, 36 miles'; 1 280 m; 20 August 1903.  
Collected by: R.S. *Pieter Faure*.
- Solariella palirrous* Barnard, 1963a: 12, pl. 2 (figs 7–8).  
Holotype: SAM-A29931. Shell broken.  
Locality: 36°48'S 52°08'E; 400 m; 9 July 1961.  
Collected by: R.S. *Africana II*.
- Solariella persculpta* Sowerby, 1903: 223, pl. 5 (fig. 8)  
= *Calliotropis persculpta* (Sowerby): Thiele, 1925: 48.  
Syntype: SAM-A5251. Shell.  
Locality: off Cape Natal; 800 m; 4 April 1901.  
Collected by: R.S. *Pieter Faure*.
- Solariella turbynei* Barnard, 1963d: 243, fig. 11a.  
Holotype: SAM-A9276. Shell.  
Locality: off Cape Point; 246 m; 17 February 1902.  
Collected by: R.S. *Pieter Faure*.
- Turcica helix* Barnard, 1964c: 21, fig. e.  
Holotype: SAM-A9295. Shell.  
Locality: off Cape Vidal; 146–183 m; 27 February 1901.

Paratype: SAM-A9256. Shell, broken.

Locality: off Umkomaas; 73 m; 31 December 1900.

Collected by: R.S. *Pieter Faure*.

*Turcica konos* Barnard, 1964c: 20, fig. 3f.

Syntypes: SAM-A9257. Four shells (two adult, one immature, one broken).

Locality: off Umkomaas River; 73 m; 31 December 1900.

Collected by: R.S. *Pieter Faure*.

*Turcica salpinx* Barnard, 1964c: 19, fig. 3a-d.

Syntype: SAM-A9252. Shell, adult, broken.

Locality: off Cape Morgan; 40 m; 26 July 1901.

Syntypes: SAM-A9253. Five shells (three broken, two juvenile).

Locality: off Hood Point; 90 m; 15 July 1901.

Collected by: R.S. *Pieter Faure*.

#### Family Liotiidae

*Brookula coronis* Barnard, 1963b: 12, pl. 2 (figs 5-6).

Syntypes: SAM-A29929. Nine shells.

Locality: 36°48'S 52°08'E; 400 m; 9 July 1961.

Collected by: R.S. *Africana II*.

*Leptothyra echomenus* Barnard, 1963b: 12, pl. 2 (figs 3-4).

Syntypes: SAM-A29928. Five shells.

Locality: 36°48'S 52°08'E; 400 m; 9 July 1961.

Collected by: R.S. *Africana II*.

*Leptothyra sola* Barnard, 1963d: 231, fig. 9e.

Holotype: SAM-A9288. Shell, probably juvenile.

Locality: probably Algoa Bay; no depth or date.

Collected by: R.S. *Pieter Faure*.

*Leptothyra syngenes* Barnard 1963b: 11, pl. 2, figs 2, 4.

Syntypes: SAM-A29927. Twelve shells, some broken.

Locality: 36°48'S 52°08'E; 400 m; 9 July 1961.

Collected by: R.S. *Africana II*.

*Rufulana sextula* Barnard, 1963d: 232, fig. 9f.

Syntypes: SAM-A29634. Three shells.

Locality: probably Algoa Bay; 55 m; 10 November 1898.

Collected by: R.S. *Pieter Faure*.



Family **Turbinidae**

*Astraliium (Cyclocantha) gilchristi* Sowerby 1903: 221, pl. 5 (fig. 6)

= *Astrea gilchristi* (Sowerby): Barnard 1963*d*: 221, fig. 7c.

Syntype: SAM-A5219. Shell.

Locality: off Scottburgh; 168 m; 7 March 1901.

Collected by: R.S. Pieter Faure.

*Calcar rhyssopoma* Barnard, 1964*b*: 17, fig. 2a-b.

Syntypes: SAM-A9285. Fourteen shells, one operculum, radula slide.

Locality: off Umkomaas River; 73 m; 31 December 1900.

Collected by: R.S. Pieter Faure.

*Turbo boswellae* Barnard, 1969: 652, fig. 27a.

Holotype: SAM-A30031. Shell, operculum, radula slide.

Locality: off Cape Point; 366 m; no date.

Collected by: commercial trawler.

*Turbo foveolatus* Barnard, 1963*d*: 216, fig. 6.

Syntypes: SAM-A9284. Five shells.

Locality: 34°05'S 25°52'E; 95 m; 14 November 1898.

Collected by: R.S. Pieter Faure.

Family **Rissoidae**

*Amphithalamus muiri* Barnard, 1963*a*: 188, fig. 15d.

Syntypes: SAM-A9262. Eleven shells.

Locality: Still Bay; intertidal zone; no date.

Collected by: T. Muir.

Family **Vitrinellidae**

*Cochliopsis catherinae* Barnard 1963*a*: 180, fig. 35a.

Holotype: SAM-A9221. Shell.

Locality: 34°27'S 25°42'E; 473 m; 14 November 1898.

Paratype: SAM-A9341. Shell.

Locality: 33°50'S 25°54'E; 61 m; 11 November 1898.

Collected by: R.S. Pieter Faure.

*Cochliopsis tugelae* Barnard, 1963*a*: 180, fig. 35b.

Holotype: SAM-A9282. Shell.

Locality: off Tugela River; 26 m; 16 January 1901.

Collected by: R.S. Pieter Faure.

*Vitrinella inclinans* Barnard 1963a: 179, fig. 35c.

Syntypes: SAM-A29680. Six shells with animals, one operculum.

Locality: 34°27'S 25°45'E; 463 m; 19 November 1898.

Collected by: R.S. Pieter Faure.

#### Family **Turritellidae**

*Turritella chrysotoxa* Tomlin, 1925: 315, fig. 3a.

Syntype: SAM-A3561. Shell.

Locality: off Illovo River; 50 m; 31 December 1900.

Collected by: R.S. Pieter Faure.

*Turritella kowiensis* Sowerby, 1900: 6, pl. 1 (fig. 12)

= *Turritella carinifera* form *kowiensis* (Sowerby): Barnard, 1963a: 166.

Syntypes: SAM-A3315. Two shells.

Locality: Kowie, Port Alfred; no depth; 1915.

Collected by: H. Becker.

*Turritella salisburyi* Tomlin, 1925: 314, fig. 36

= *Turritella sanguinea* Reeve: Barnard, 1963a: 169.

Syntype: SAM-A5113. Shell.

Syntype: SAM-A33952. Shell.

Locality: off East London; 95 m; 12 July 1901.

Collected by: R.S. Pieter Faure.

#### Family **Vermetidae**

*Vermetus (Stoa) corralinaceus* Tomlin, 1939: 145, pl. 12 (fig. 4).

Paratypes: SAM-A31494. Two blocks of tubes, radula slide.

Locality: Oudekraal, west Cape Peninsula; no depth or date.

Collected by: University of Cape Town.

*Vermetus periscopium* Barnard, 1963a: 146, fig. 30a.

Paratypes: SAM-A29925. Four shells, one operculum, four animals, radula slide.

Locality: 29°46'S 31°17'E; 110–130 m; 23 April 1958.

Collected by: University of Cape Town.

*Vermetus quincunx* Barnard, 1963a: 148, figs 28c, 29b–c.

Syntypes: SAM-A29814. Large mass, numerous fragments, protoconchs, two opercula, radula slide.

Locality: 34°27'S 25°42'E; 470 m; 19 November 1898.

Collected by: R.S. Pieter Faure.

Family **Cerithiidae**

*Alabina telamon* Barnard, 1963a: 138, fig. 26d.

Holotype: SAM-A29676. Shell.

Locality: off Tugela River; 118–146 m; 11 January 1901.

Collected by: R.S. Pieter Faure.

Family **Cerithiopsidae**

*Cerithiella natalensis* Barnard, 1963a: 126.

Holotype: SAM-A9045. Shell.

Locality: off Umkomaas River; 73 m; 31 December 1900.

Collected by: R.S. Pieter Faure.

*Cerithiella nonnitens* Barnard, 1963a: 127, fig. 23g.

Holotype: SAM-A9047. Shell.

Locality: off Cove Rock, East London; 146–183 m; 30 July 1900.

Collected by: R.S. Pieter Faure.

*Cerithiella (Stilus) subuliapex* Barnard, 1963b: 10, pl. 1 (fig. 8).

Holotype: SAM-A29924. Shell.

Locality: 27°48'S 47°19'E; 875 m; 22 June 1961.

Collected by: R.S. Africana II.

*Cerithiella taylori* Barnard, 1963a: 126, fig. 23i.

Holotype: SAM-A9711. Shell.

Locality: 33°26'S 16°33'E; 2 260–2 370 m; August 1959.

Collected by: R.S. Africana II.

*Cerithiella vidalensis* Barnard, 1963a: 127.

Holotype: SAM-A9046. Shell.

Locality: off Cape Vidal; 146–183 m; 27 February 1901.

Collected by: R.S. Pieter Faure.

*Cerithiopsis rota* Barnard, 1963a: 124, fig. 23a–c.

Holotype: SAM-A9043. Shell.

Locality: 34°25'S 44°33'E; 61 m; 10 November 1898.

Collected by: R.S. Pieter Faure.

Family **Finellidae**

*Cerithidium fragrans* Barnard, 1963a: 139, fig. 15c.

Syntypes: SAM-A29679. Five shells, fossils.

Locality: raised beach, Little Brak River; no date.

Collector unknown.

Family *Abyssochrysidae*

*Abyssochrysos tomlini* Barnard, 1963a: 143, fig. 27f

= *Abyssochrysos melvilli* (Schepman): Houbbrick, 1979: 10, figs 7d, f, i, 8.

Lectotype: SAM-A9101. Shell.

Paralectotype: SAM-A33950 (previously A9109). Shell.

Locality: off Cape Natal; 535 m; 4 April 1901.

Collected by: R.S. Pieter Faure.

*Abyssochrysos melanoides* Tomlin, 1927: 78. figs 1-3.

Syntype: SAM-A5116. Shell.

Locality: off Cape Point; 1 462-2 724 m; 21 July 1903.

Syntypes: SAM-A33951 (previously A5116). Twelve shells, four opercula, one radula.

Collected by: R.S. Pieter Faure.

Family *Triforidae*

*Trifora ? axialis* Barnard, 1963b: 10, pl. 1 (fig. 7) (*sic*).

Syntypes: SAM-A29923. Two shells.

Locality: 36°48'S 52°08'E; 400 m; 9 July 1961.

Collected by: R.S. *Africana II*.

*Trifora bactron* Barnard, 1963a: 115, fig. 20c.

Holotype: SAM-A9077. Shell.

Locality: off Umkomaas River; 73 m; 31 December 1900.

Paratype: SAM-A9076. Shell.

Locality: off Umhloti River; 73 m; 18 December 1900.

Collected by: R.S. Pieter Faure.

*Trifora baculus* Barnard, 1963a: 115, fig. 20b.

Holotype: SAM-A9078. Shell.

Locality: off Sandy Point north of Cape Morgan; 93 m; 14 August 1901.

Collected by: R.S. Pieter Faure.

*Trifora dagama* Barnard, 1963a: 112, fig. 19f.

Syntype: SAM-A9073. Shell.

Locality: off Umkomaas River; 73 m; 31 December 1900.

Collected by: R.S. Pieter Faure.

*Trifora morgani* Barnard, 1963a: 111, fig. 19a.

Holotype: SAM-A9079. Shell, broken.

Locality: off Sandy Point north of Cape Morgan; 93 m; 14 August 1901.

Collected by: R.S. Pieter Faure.

*Trifora natalensis* Barnard, 1963a: 112, fig. 19c.

Syntype: SAM-A9071. Shell.

Locality: off Umhloti River; 73 m; 18 December 1900.

Syntype: SAM-A9072. Shell.

Locality: off Umkomaas River; 73 m; 31 December 1900.

Collected by: R.S. Pieter Faure.

*Trifora scala* Barnard, 1963a: 115, fig. 19d.

Syntype: SAM-A9074. Shell.

Locality: off Umkomaas River; 73 m; 31 December 1900.

Syntype: SAM-A9075. Shell.

Locality: off Umhloti River; 73 m; 18 December 1900.

Collected by: R.S. Pieter Faure.

#### Family Pyramidellidae

*Odostomia tugelae* Barnard, 1963a: 93, fig. 15b.

Holotype: SAM-A9225. Shell.

Locality: off Tugela River; 65–80 fm.; 11 January 1901.

Collected by: R.S. Pieter Faure.

*Turbonilla bathybias* Barnard, 1963a: 89.

Syntypes: SAM-A9175. One shell, one apex.

Locality: off East London; 400–450 fm.; 17 April 1901.

Collected by: R.S. Pieter Faure.

*Turbonilla? biflosa* Barnard, 1963a: 90, fig. 15a.

Holotype: SAM-A9220. Shell.

Locality: 34°05'S 25°55'E; 122 m; 1 November 1898.

Collected by: R.S. Pieter Faure.

*Turbonilla deprofundis* Barnard, 1963a: 89, fig. 14b.

Holotype: SAM-A9157. Shell.

Locality: off East London; 400–450 fm.; 17 April 1901.

Collected by: R.S. Pieter Faure.

*Turbonilla tugelae* Barnard, 1963a: 88, fig. 14f.

Syntypes: SAM-A9153. Shell, apex.

Locality: off Tugela River; 14 fm.; 16 January 1901.

Collected by: R.S. Pieter Faure.

#### Family Epitoniidae

*Acrilla analogica* Barnard, 1963a: 106, fig. 18g–h.

Holotype: SAM-A9026. Protoconch with nine whorls.

Locality: 36°40'S 21°26'E; 366 m; 17 July 1906.

Paratype: SAM-A9229. Shell, broken.  
Locality: 34°27'S 25°42'E; 467 m; 14 November 1898.  
Collected by: R.S. Pieter Faure.

*Acrilla natalis* Barnard, 1963a: 105, fig. 18e-f.  
Syntypes: SAM-A9028. Two shells, broken.  
Locality: off Cape Natal; 99 m; 14 December 1900.  
Collected by: R.S. Pieter Faure.

*Scala blaisei* Barnard, 1963a: 102, fig. 18a  
= *Epitonium blaisei* (Barnard): **comb. nov.**  
Holotype: SAM-A9025. Shell.  
Locality: off Cape St. Blaize; 228 m; 21 December 1899.  
Collected by: R.S. Pieter Faure.

*Scala bonaspei* Barnard, 1963a: 104, fig. 17b  
= *Epitonium bonaspei* (Barnard): **comb. nov.**  
Syntypes: SAM-A9761. Two shells with opercula, proboscis, jaws, radula  
of larger specimen on slide.  
Syntype: SAM-A9793. Shell.  
Locality: off Cape Point; 3 120-3 260 m; December 1959.  
Collected by: R.S. *Africana II*.

*Scala illovoensis* Barnard, 1963a: 101, fig. 18b  
= *Epitonium illovoensis* (Barnard): **comb. nov.**  
Holotype: SAM-A9023. Shell.  
Locality: off Illovo River; 50-55 m; 31 December 1900.  
Collected by: R.S. Pieter Faure.

*Scala mathildopsis* Barnard, 1963b: 9, pl. 1 (fig. 5)  
= *Epitonium mathildopsis* (Barnard): **comb. nov.**  
Syntypes: SAM-A29921. Six shells.  
Locality: 27°48'S 47°19'E; 875 m; 22 June 1961.  
Collected by: R.S. *Africana II*.

*Scala munda* Barnard, 1969: 649, fig. 24b  
= *Epitonium mundum* (Barnard): **comb. nov.**  
Holotype: SAM-A30552. Shell.  
Locality: off Cape Point; 1 250 m; 20 August 1903.  
Paratypes: SAM-A7419. Three shells.  
Locality: off Cape Point; 1 280 m; 20 August 1903.  
Collected by: R.S. Pieter Faure.

*Scala tenebrosa* Sowerby, 1903: 220, pl. 4 (fig. 6)

= *Epitonium tenebrosum* (Sowerby): **comb. nov.**

Syntype: SAM-A3599. Shell.

Locality: off Cape St. Blaize; 67 m; 22 October 1900.

Collected by: R.S. Pieter Faure.

#### Family Mathildidae

*Mathilda salve* Barnard, 1963a: 176, fig. 34.

Holotype: SAM-A9139. Shell.

Locality: off Cape St. Blaize; 229 m; 21 December 1899.

Paratype: SAM-A9141. Apex of seven whorls, no protoconch.

Locality: off Glendower Beacon, Port Alfred; 183 m; 26 July 1901.

Collected by: R.S. Pieter Faure.

#### Family Architectonicidae

*Fluxina solarium* Barnard, 1963b: 11, pl. 2 (fig. 1).

Holotype: SAM-A29926. Shell.

Locality: 27°48'S 47°19'E; 875 m; 22 June 1961.

Collected by: R.S. *Africana II*.

*Heliacus numulus* Barnard, 1963a: 163, fig. 31e.

Holotype: SAM-A9125. Shell.

Locality: off Cape Morgan; 141 m; 26 July 1901.

Collected by: R.S. Pieter Faure.

*Heliacus obolos* Barnard, 1963a: 163, fig. 31f-g.

Holotype: SAM-A9127. Shell, broken.

Locality: off Umhloti River; 73 m; 31 December 1900.

Collected by: R.S. Pieter Faure.

*Heliacus petasus* Tomlin, 1928: 334, pl. 26 (fig. 4).

Holotype: SAM-A3579. Shell.

Locality: off Scottburgh; 169 m; 7 March 1901.

Collected by: R.S. Pieter Faure.

#### Family Stiliferidae

*Apicalia sowerbyi* Barnard, 1963a: 94, fig. 16.

Syntypes: SAM-A9169. Two shells.

Locality: off Hood Point, near East London; 90 m; 15 July 1901.

Collected by: R.S. Pieter Faure.

Family **Fossaridae**

*Fossarus translucens* Barnard, 1969: 647, fig. 23a.

Holotype: SAM-A30052. Shell.

Locality: Gordon's Bay, False Bay; intertidal; no date.

Collected by: Dr Ackert.

*Megalomphalus mosselensis* Barnard, 1957: 180, figs b-d.

Holotype: SAM-A29922. Shell, six embryo ex utera, radula slide.

Locality: Mossel Bay; no depth or date.

Collected by: University of Cape Town.

Family **Amaltheidae**

*Cheila microstriata* Barnard, 1963a: 76, fig. 10j.

Syntype: SAM-A9004. Shell, broken.

Locality: off Durban; 153 m; 17 December 1900.

Syntype: SAM-A9005. Shell.

Locality: off Umhloti River; 73 m; 18 December 1900.

Syntype: SAM-A9006. Shell, broken.

Locality: off Durnford Point, Natal; 164 m; 11 July 1900.

Syntype: SAM-A9342. Shell.

Locality: off Cape Vidal; 146-183 m; 27 February 1901.

Collected by: R.S. Pieter Faure.

Family **Trichotropidae**

*Trichotropis zuluensis* Barnard, 1963a: 78, figs 12d-13c.

Syntype: SAM-A9011. Shell, radula slide.

Locality: off Durnford Point, Natal; 165 m; 28 February 1901.

Syntype: SAM-A9012. Shell.

Locality: off Cape Vidal; 146-183 m; 27 February 1901.

Collected by: R.S. Pieter Faure.

Family **Aporrhaidae**

*Aporrhais pesgallinae* Barnard, 1963a: 67, fig. 8.

Holotype: SAM-A30248. Shell with animal in alcohol, radula slide.

Locality: 26°23'S 14°17'E; 311 m; 28 October 1948.

Collected by: R.S. *Africana I.*

Family **Naticidae**

*Falsilunatia pseudopsila* Barnard, 1963a: 64, fig. 7e; 1969: 646, fig. 22.

Holotype: SAM-A9780. Shell with operculum, radula slide.



Locality: off Cape Point; 2 920 m; December 1959.

Collected by: R.S. *Africana II*.

*Polynices cleistopsila* Barnard, 1963a: 64, fig. 7d.

Syntypes: SAM-A9735. One adult shell, two juveniles, three opercula, radula slide.

Locality: Off Cape Point, 33°36'S 16°15'E; 2 830 m; August 1959.

Collected by: R.S. *Africana II*.

#### Family Cypraeidae

*Cypraea fultoni* Sowerby, 1903: 218, pl. 4 (fig. 7).

Holotype: SAM-A5053. Shell.

Locality: South Africa; *ex pisce*; no depth or date.

Collected by: R.S. *Pieter Faure*.

#### Family Triviidae

*Trivia verhoefi* Gosliner & Liltved, 1982: 121, figs 1f, 13-16.

Holotype: SAM-A34387. Shell with animal.

Locality: Llandudno, 34°01'S 18°20'E; 25 m; 22 December 1979.

Collected by: G. Verhoef.

Paratype: SAM-A34388. Shell with animal.

Locality: Llandudno; 20 m; 16 March 1980.

Collected by: W. R. Liltved.

Paratype: SAM-A34389. Shell with animal.

Locality: Llandudno, 25 m; 27 March 1980.

Collected by: W. R. Liltved.

#### Family Ovulidae

*Amphiperas beckeri* Sowerby, 1900: 5, pl. 1 (fig. 13)

= *Primovula beckeri* (Sowerby): Thiele, 1929; Barnard 1963a: 55, figs 5d-6c.

Paratypes: SAM-A8918. Four shells.

Locality: Kowie, Port Alfred; intertidal; ?1899.

Collected by: H. Becker.

#### Family Tonnidae

*Eudolium aulacoides* Tomlin, 1927: 83, fig. 4a.

Holotype: SAM-A3529. Shell.

Locality: off Cape Point; 1 020-1 280 m; 17 September 1903.

Collected by: R.S. *Pieter Faure*.

## Family Cymatiidae

*Thalassocyon bonus* Barnard, 1960: 440, fig. 3.

Holotype: SAM-A9714. Shell, operculum broken, radula slide A9810.

Locality: off Cape Point; 2 280-2 380 m; August 1959.

Collected by: R.S. *Africana II*.

## Family Columbariidae

*Columbarium angulare* Barnard, 1959: 235, fig. 52c.

Syntypes: SAM-A4608. Three shells, operculum, radula slide.

Locality: off Cape Point; 1 480-1 650 m; 14 July 1903.

Collected by: R.S. *Pieter Faure*.

*Columbarium formosissimum* Tomlin, 1928: 331, pl. 25 (fig. 1).

Holotype: SAM-A3501. Shell.

Locality: Cape St. Blaize; 'N by E  $\frac{1}{4}$  E 65 miles'; 90 fm.; 22 December 1899.

Collected by: R.S. *Pieter Faure*.

Paratype: SAM-A3499; Shell.

Locality: Cape Seal; 'N by W  $\frac{1}{2}$  W 55 miles'; 87 fm.; 19 February 1902.

Collected by: R.S. *Pieter Faure*.

Paratype: SAM-A3450; Shell.

Locality: Cape Seal; 'N by W  $\frac{1}{2}$  W 55 miles'; 87 fm.; 19 February 1902.

Collected by: R. S. *Pieter Faure*.

Paratype: SAM-A3498; Shell.

Locality: Glendower Beacon; 'N  $\frac{1}{2}$  W 6 miles'; 66 fm.; 10 September 1901.

Collected by: R.S. *Pieter Faure*.

*Columbarium natalense* Tomlin, 1928: 331, pl. 25 (fig. 2).

Holotype: SAM-A3497. Shell.

Locality: off Cape Natal; 153 m; 17 December 1900.

Collected by: R.S. *Pieter Faure*.

*Columbarium rotundum* Barnard, 1959: 236, fig. 52d.

Syntypes: SAM-A4592. Two shells, two opercula.

Locality: off Cape Point; 1 400 m; 27 August 1903.

Collected by: R.S. *Pieter Faure*.

## Family Colubrariidae

*Epidromus crebrilirata* Sowerby, 1903: 220, pl. 4 (fig. 4)

= *Nivitriton fictilis* (Hinds): Kilburn, 1977: 194.

Holotype: SAM-A4974. Shell.

Locality: off Glendower Beacon, Port Alfred; 183 m; 10 September 1901.

Collected by: R.S. *Pieter Faure*.

Family **Muricidae**

*Tritonalia juritzi* Barnard, 1969: 639, fig. 18d.

Holotype: SAM-A30051. Shell, broken.

Locality: doubtful, probably Cape Peninsula.

Collected by: C. Juritz.

*Trophon acceptans* Barnard, 1959: 202, figs 40d-43b; 1963c; 435, fig. 7a; 1969: 638.

Holotype: SAM-A3473. Shell with operculum, radula slide.

Locality: off Cape Point, 'N 64° E 37 miles'; 1 270-1 460 m; 9 September 1903.

Paratype: SAM-A3449. Three shells.

Locality: off Cape Point, 'E ¾° N 38 miles'; 1 190 m; 9 September 1903.

Paratype: SAM-A3480. Shell.

Locality: off Cape Point, '64° NE 37 miles'; 1 270-1 460 m; 10 September 1903.

Collected by: R.S. Pieter Faure.

*Trophon? incertus* Barnard, 1959: 203, fig. 32a.

Syntypes: SAM-A8843. Two shells.

Locality: off Cape Natal; 800 m; 4 April 1901.

Collected by: R.S. Pieter Faure.

*Trophon johanthielei* Barnard, 1959: 206, fig. 44b.

Holotype: SAM-A8844. Shell, broken.

Locality: off East London; 730-820 m; 17 April 1901.

Collected by: R.S. Pieter Faure.

*Trophon mioplectos* Barnard, 1959: 206, fig. 44a.

Holotype: SAM-A8631. Shell.

Locality: 34°27'S 25°42'E; 466 m; 16 November 1898.

Collected by: R.S. Pieter Faure.

*Trophon pistillum* Barnard, 1959: 206, fig. 44f.

Holotype: SAM-A8842. Shell, broken; radula slide.

Locality: off Cape Natal; 802 m; 4 April 1901.

Collected by: R.S. Pieter Faure.

*Typhus pentaphasios* Barnard, 1959: 211, fig. 46h-i

= *Siphonochelus pentaphasios* (Barnard 1959): Radwin & D'Attilo 1976: 199, fig. 144.

Syntypes: SAM-A4949. Two shells.

Locality: off Cape Point; 1 204 m; 10 September 1903.

Collected by: R.S. Pieter Faure.

Family **Coralliophilidae**

*Coralliophila isosceles* Barnard, 1959: 191, fig. 39a.

Syntypes: SAM-A4951. Two shells.

Locality: off Glendower Beacon, Port Alfred; 121 m; 10 September 1901.

Collected by: R.S. Pieter Faure.

*Coralliophila zuluensis* Barnard, 1959: 192, fig. 39b.

Holotype: SAM-A8852. Shell.

Locality: off O'Neil Peak, Natal; 163 m; 28 February 1901.

Collected by: R.S. Pieter Faure.

*Latiaxis capensis* Tomlin, 1928: 332.

Holotype: SAM-A3522. Shell.

Locality: off Sandy Point, Transkei; 173 m; 14 August 1901.

Collected by: R.S. Pieter Faure.

*Latiaxis? elstoni* Barnard, 1962b: 148, fig. 1.

Holotype: SAM-A9346. Shell.

Locality: Natal coast; *ex pisce*; donated 1961.

Collector unknown.

*Latiaxis kyllix* Barnard, 1959: 188, fig. 27

= *Latiaxis mawae kyllix* Barnard: Kilburn, 1973: 565, fig. 9b.

Holotype: SAM-A8850. Shell, juvenile.

Locality: off Cape Natal; 98 m; 14 December 1900.

Collected by: R.S. Pieter Faure.

Family **Buccinidae**

*Babylonia pintado* Kilburn, 1971: 486, figs 3, 7c, e.

Paratypes: SAM-A4751. Shell.

Locality: off Port Shepstone, 'WNW 8 miles'; 27 fm.; 14 March 1901.

Collected by: R.S. Pieter Faure.

Paratype: SAM-A7706. Shell.

Locality: Natal; *ex pisce*; no depth or date.

Collected by: Mr Casey.

Paratype: SAM-A8239. Shell.

Locality: Natal; *ex pisce*; no depth or date.

Collected by: H. W. Bell-Marley.

Paratypes: SAM-A30956. Six shells.

Locality: Natal; *ex pisce*; no depth or date.

Collected by: R. Frames.

*Glypteuthria solidissima* Tomlin, 1932: 166, fig. 7

= *Afrocominella elongata* (Dunker): Barnard, 1959: 153, fig. 31f.

Holotype: SAM-A3543. Shell.

Locality: off Cape Point; 82 m; 6 June 1900.

Collected by: R.S. Pieter Faure.

*Neptunea bonaespei* Barnard, 1960: 440, fig. 3.

Syntypes: SAM-A9826. Three shells (one broken), operculum, radula slide.

Locality: off Cape Point, 33°52'S 16°51'E; 2 520–2 820 m; December 1959.

Collected by: R.S. *Africana II*.

*Nux alabaster* Barnard, 1960: 440, fig. 3.

Holotype: SAM-A9715. Shell with operculum, radula slide.

Locality: off Cape Point, 33°26'S 16°33'E; 2 270–2 380 m; December 1959.

Paratypes: SAM-A9827. Two shells with opercula.

Locality: off Cape Point; 35°52'S 16°51'E; 2 540–2 820 m; December 1959.

Collected by: R.S. *Africana II*.

*Prosipho torquatus* Barnard, 1963c: 433, fig. 6c–e.

Holotype: SAM-A9884. Shell with operculum, radula slide.

Locality: off Cape Point, 34°37'S 17°03'E; 2 890–2 960 m; December 1959.

Collected by: R.S. *Africana II*.

#### Family Columbellidae

*Columbella beckeri* Sowerby, 1900: 3, pl. 1 (fig. 7).

Syntype: SAM-A3322. Shell.

Locality: Port Alfred; intertidal; 1915.

Collected by: H. Becker.

*Columbella confertilirata* Barnard, 1959: 182, fig. 35d.

Syntypes: SAM-A8879. One adult, two juveniles, fifteen fragments.

Locality: off Umkomaas; 73 m; 31 December 1900.

Collected by: R.S. Pieter Faure.

*Columbella dibolos* Barnard, 1964c: 17

= *Mitrolumna stepheni* (Melvill & Standen): Kilburn, 1977: 204.

Syntypes: SAM-A29636. Three worn juveniles, one broken.

Locality: off Umhloti River; 73 m; 18 December 1900.

Collected by: R.S. Pieter Faure.

*Columbella lightfooti* Smith, 1901; 112, pl. 1 (fig. 3).

Paratypes: SAM-A6936. Three shells.

Locality: Kalk Bay, False Bay; no depth or date.

Collected by: R. M. Lightfoot.

'*Columbella*' *polyarosus* Barnard, 1959: 182, fig. 35c.

Holotype: SAM-A8875. Shell, broken.

Locality: off Cape Natal; 99 m; 14 December 1900.

Paratype: SAM-A8876. Apex.

Locality: off O'Neil Peak; 165 m; 28 February 1901.

Paratypes: SAM-A8877. Shell, three fragments.

Locality: off Umhloti River; 73 m; 18 December 1900.

Paratype: SAM-A8878. Apex.

Locality: off Hood Point; 90 m; 15 July 1901.

Collected by: R.S. Pieter Faure.

'*Columbella*' *sigma* Barnard, 1959: 185, fig. 36b.

Holotype: SAM-A8890. Shell.

Locality: off Cove Rock, East London; 40 m; 6 August 1901.

Collected by: R.S. Pieter Faure.

'*Columbella*' *vitula* Barnard, 1959: 184, fig. 36a.

Holotype: SAM-A8887. Shell, juvenile.

Locality: off Cove Rock, East London; 80–130 fm.; 30 July 1901.

Collected by: R.S. Pieter Faure.

*Pyrene parhelena* Barnard, 1959: 177, fig. 36b.

Syntypes: SAM-A8870. Two shells (one broken), radula slide.

Locality: off Cape St. Blaize; 229 m; 21 December 1899.

Collected by: R.S. Pieter Faure.

*Pyrene salmoneus* Barnard, 1963b: 7.

Syntypes: SAM-A29920. One hundred and fifty-six shells, opercula, many fragments.

Locality: South of Madagascar, 36°48'S 52°08'E; 400 m; 9 July 1961.

Collected by: R.S. *Africana II*.

#### Family Nassariidae

*Nassa mui* Barnard, 1959: 105.

Syntypes: SAM-A29919. Eleven shells.

Locality: uncertain.

Collected by: University of Cape Town.

*Nassa rhysonopia* Barnard, 1969: 627, fig. 15a.

Holotype: SAM-A29990. Shell, juvenile.

Paratype: SAM-A30551. Shell, juvenile.

Locality: off Cove Rock, East London; 40 m; 6 August 1901.

Collected by: R.S. Pieter Faure.

*Nassa scopularcus* Barnard, 1959: 120, fig. 24a.

Holotype: SAM-A9230. Shell.

Locality: Late Tertiary marine beds, lagoon shore, south of Bogenfels, South West Africa; 1930.

Collected by: S. H. Haughton, University of Cape Town.

*Nassa vidalensis* Barnard, 1959: 118, fig. 24b.

Holotype: SAM-A8839. Shell.

Locality: off Cape Vidal; 146–183 m; 27 February 1901.

Collected by: R.S. Pieter Faure.

#### Family Fasciolariidae

*Fasciolaria agulhasensis* Tomlin, 1932: 157, fig. 1

= *Fasciolaria (Pleuroploca) scholveni* Strebel: Kilburn, 1974: 204, fig. 13a.

Holotype: SAM-A6539. Shell.

Locality: Agulhas Bank; 28 fm.; 5 January 1904.

Collected by: R.S. Pieter Faure.

*Fasciolaria? holcophorus* Barnard, 1959: 98, fig. 21.

Holotype: SAM-A8819. Shell.

Locality: off Cape St. Blaize; 228 m; 21 December 1899.

Collected by: R.S. Pieter Faure.

*Fusus africanae* Barnard, 1959: 96, figs 19k–20d.

Holotype: SAM-A8610. Shell, juvenile with operculum, radula slide.

Locality: Brown's Bank; 146–183 m; 17 July 1906.

Paratype: SAM-A8826. Shell, juvenile, radula slide.

Locality: off western Cape Peninsula; 256–276 m; 28 March 1900.

Collected by: R.S. Pieter Faure.

*Fusus bonaspei* Barnard, 1959: 95, fig. 20c–f.

Holotype: SAM-A4632. Shell with operculum.

Locality: doubtful, wrong *Pieter Faure* station number recorded but probably off Cape Point, depth and date unknown.

Paratypes: SAM-A4629. Two shells with opercula.

Locality: off Cape Point, 'NE  $\frac{1}{2}$  N 19 miles'; 266 m; 27 February 1902.

Paratype: SAM-A4630. Six shells, one operculum.

Locality: off Cape Hangklip 'NE  $\frac{3}{4}$  N 28 miles'; 183 m; 27 August 1903.

Collected by: R.S. Pieter Faure.

*Fusus faurei* Barnard, 1959: 94, figs 19h–20e.

Syntypes: SAM-A4581. Nine shells, six opercula, radula slide.

Locality: off Cape Point, 'N 41° E 38 miles'; 578–730 m; 16 September 1903.

Syntype: SAM-A4582. Shell.

Locality: off Cape Point, 'NE  $\frac{1}{2}$  N 39 miles'; 580–1 020 m; 17 September 1903.

Collected by: R.S. *Pieter Faure*.

*Latirus mosselensis* Tomlin, 1932: 158, fig. 1.

Syntype: SAM-A3504. Shell.

Locality: off Mossel Bay; 49 m; 14 June 1899.

Collected by: R.S. *Pieter Faure*.

#### Family Volutidae

*Fulgoraria blaizei* Barnard, 1959: 28, fig. 8b; 1969: 620, fig. 11

= *Fusivoluta blaizei* (Barnard, 1959): Weaver, 1963: 5.

Holotype: SAM-A3433. Shell with operculum.

Locality: off Cape St. Blaize, 'NE  $\frac{3}{4}$  E 73 miles'; 192 m; 21 February 1902.

Paratypes: SAM-A3430. Shell, adult, 2 juveniles.

Locality: off Cape St. Blaize, 'N by E 73 miles'; 230 m; 21 December 1889.

Collected by: R.S. *Pieter Faure*.

*Glypteuthria capensis* Tomlin, 1932: 165; 1945: 135

= *Fusivoluta sculpturata* (Tomlin): Weaver & Dupont, 1970: 185.

Holotype: SAM-A3446. Shell with operculum.

Locality: off Cape Point, 'N 41° E 38 miles'; 593–730 m; 16 September 1903.

Paratype: SAM-A3448. Shell.

Locality: off Cape Point, 'NE by E  $\frac{1}{4}$  S 28 miles'; 393–733 m; 28 September 1903.

Collected by: R.S. *Pieter Faure*.

*Fusivoluta decussata* Barnard, 1959: 31, fig. 8c.

Holotype: SAM-A3432. Shell.

Locality: off East London, Buffalo River, 'N 15 miles'; 566 m; 24 April 1901.

Collected by: R.S. *Pieter Faure*.

*Fusivoluta elegans* Barnard, 1959: 32, fig. 8a.

Holotype: SAM-A8803. Shell.

Locality: off East London, Buffalo River; 830 m; 17 April 1901.

Collected by: R.S. *Pieter Faure*.

*Volutocorbis boswellae* Rehder, 1969: 202, pl. 40 (fig. 5), pl. 41 (figs 16–19).

Paratype: SAM-A2010. Shell.

Locality: off Cape St. Blaize, 'N by E  $\frac{1}{4}$  E 65 miles'; 155 m; 28 December 1899.

Paratype: SAM-A2011. Shell.



Locality: off Cape St. Blaize, 'N by E  $\frac{1}{4}$  E 67 miles'; 164 m; 22 December 1899.

Paratype: SAM-A2009. Shell.

Locality: off Cape Seal, 'N by E  $\frac{3}{4}$  E 37 miles'; 146 m; 20 February 1902.

Paratypes: SAM-A31976 (previously SAM-A3335). Six shells.

Locality: Brown's Bank, 36°40'S 21°26'E; 146–183 m; 17 July 1906.

Collected by: R.S. Pieter Faure.

*Volutocorbis dispirilis* Rehder, 1969: 203, pl. 40 (fig. 4), pl. 41.

Holotype: SAM-A31977 (previously SAM-A3335). Shell.

Paratype: SAM-A33687 (previously SAM-A3335). Shell.

Locality: Brown's Bank, 36°40'S 21°26'E; 146–183 m; 17 July 1906.

Collected by: R.S. Pieter Faure.

*Volutilithes gilchristi* Sowerby, 1902: 99, pl. 2 (fig. 5)

= *Volutocorbis gilchristi* (Sowerby): Barnard, 1959: 28.

Paratype: SAM-A1974. Shell.

Locality: off Cape Natal, 'W by N  $\frac{3}{4}$  N 11 miles'; 189–366 m; 1 April 1901.

Collected by: R.S. Pieter Faure.

#### Family Marginellidae

*Diliculum inopinatum* Barnard, 1962a: 14, fig. 1.

Paratypes: SAM-A29917. Three shells, radula slide.

Locality: Saldanha Bay; 3–28 fm.; 5 May 1960.

Collected by: University of Cape Town.

*Persicula alborubida* Barnard, 1969: 620, fig. 10d.

Holotype: SAM-A30050. Shell.

Locality: Kommetjie; intertidal; no date.

Collected by: C. Connolly.

Paratype: SAM-A30550. Shell.

Locality: Kommetjie; intertidal; no date.

Collected by: C. Connolly.

*Persicula nigrocrocea* Barnard, 1969: 619, fig. 10b–c.

Holotype: SAM-A30049. Shell.

Locality: Simonstown; no depth or date.

Collected by: C. Connolly.

Paratype: SAM-A30549. Shell.

Locality: Simonstown; no depth or date.

Collected by: C. Connolly.

*Marginella brocktoni* Shackleford, 1914: 98, two figures.

Syntypes: SAM-A1956. Shell with animal, one shell in pieces.

Locality: off Cape Point, 'N 50° E 18 miles'; 329 m; 27 February 1902.  
Collected by: R.S. Pieter Faure.

*Marginella differens eugenes* Barnard, 1959: 8; 1969: 619, fig. 10a.

Holotype: SAM-A8786. Shell.

Locality: off Umkomaas River; 73 m; 31 December 1900.

Collected by: R.S. Pieter Faure.

*Marginella kerochuta* Shackleford, 1914: 97, two figures.

Syntypes: SAM-A1954. Two shells.

Locality: off Cape Point, 'NE  $\frac{1}{4}$  N 18 miles'; 244 m; 27 February 1902.

Collected by: R.S. Pieter Faure.

*Marginella taylori* Shackleford, 1916: 194, figs 3-4

= *Marginella differens* Smith: Barnard, 1959: 8.

Syntypes: SAM-A7191. Two shells, one broken.

Locality: off Cape St. Blaize, 'N by E  $\frac{1}{2}$  E'; 193 m; no date.

Collected by: R.S. Pieter Faure.

*Marginella tomlini* Shackleford, 1916: 193, figs 1-2

= *Marginella bicatenata* Sowerby: Barnard, 1959: 6.

Holotype: SAM-A3704. Shell with animal.

Locality: off Cape St. Blaize, 'N by E  $\frac{1}{2}$  E 68 miles'; 193 m; 21 February 1902.

Collected by: R.S. Pieter Faure.

*Marginella walvisiana* Tomlin, 1920: 88, fig. 3.

Syntypes: SAM-A30666. Three shells.

Locality: Walvis Bay; no depth or date.

Collected by: P. R. Frames.

#### Family Olividae

*Ancilla errorum* Tomlin, 1921: 216, pl. 8, fig. 2.

Holotype: SAM-A1811. Shell.

Locality: Walker Point, 'NE by N  $\frac{1}{2}$  N 7 miles'; 47 fm.; 11 October 1900.

Paratype: SAM-A1816. Shell.

Locality: off Cape Point; 42 fm.; 18 December 1900.

Collected by: R.S. Pieter Faure.

*Ancilla rouillardi* Kilburn, 1981: 381, figs 6, 50, 87, 88, 99, 104-105.

Paratype: SAM-A1797. Shell.

Locality: ?Cape St. Blaize; 40 fm.; 13 December 1900.

Collected by: R.S. Pieter Faure.

Family **Mitridae**

*Charitodoron aglaia* Tomlin, 1932: 169, fig. 9

= *Charitodoron agulhasensis* (Thiele): Barnard, 1959: 146.

Holotype: SAM-A30719 (previously A3440). Shell.

Paratype: SAM-A3440. Shell.

Locality: uncertain, no *Pieter Faure* station number recorded.

Collected by: R.S. *Pieter Faure*.

*Charitodoron euphrosyne* Tomlin, 1932: 167, fig. 8.

Holotype: SAM-A3441. Shell, broken.

Locality: off Cape Point, N 77° E; 1 220–1 275 m; 28 March 1900.

Collected by: R.S. *Pieter Faure*.

*Charitodoron thalia* Tomlin, 1932: 169, fig. 10.

Holotype: SAM-A8602 (previously A1742). Shell, radula slide.

Locality: off Cape Point; 240 m; 28 March 1900.

Collected by: R.S. *Pieter Faure*.

*Mitra (Dibaphus) bathybius* Barnard, 1959: 51, figs 11a–13b.

Holotype: SAM-A8827. Shell, broken, radula slide.

Locality: off Cape Natal; 805 m; 4 April 1901.

Collected by: R.S. *Pieter Faure*.

*Mitra (Mitra) boswellae* Cate, 1964: 219, pl. 28 (fig. 1a–b)

= *Mitra guttata* Swainson: Cernohorsky, 1976: 326.

Holotype: SAM-A29799. Shell.

Locality: uncertain; no date.

Collected by: probably commercial trawler.

*Mitra fidis* Sowerby, 1916: 491, text figure

= *Pusia patula* (Reeve): Barnard, 1958: 58.

Syntypes: SAM-A29918. Two shells.

Locality: 'South Africa'; no date.

Donated by: A. E. Salisbury, 1957.

Family **Cancellariidae**

*Admete decapensis* Barnard, 1960: 439, fig. 1a.

Holotype: SAM-A9777. Shell, partly corroded.

Locality: off Cape Point, 34°37'S 17°03'E; 3 070–3 150 m; December 1959.

Collected by: R.S. *Africana II*.

*Cancellaria euetrios* Barnard, 1959: 14, fig. 3.

Holotype: SAM-A8747. Shell.

Locality: off Cape Recife, 34°26'S 25°42'E; 227 m; 14 November 1898.  
Collected by: R.S. *Pieter Faure*.

*Cancellaria euthymei* Barnard, 1960: 438, fig. 16.

Holotype: SAM-A9888. Shell.

Locality: off Cape Point, 34°26'S 17°00'E; 2 750–3 260 m; December 1959.  
Collected by: R.S. *Africana II*.

*Cancellaria producta* Sowerby, 1903: 220, pl. 4 (fig. 5).

Syntype: SAM-A339. Shell.

Locality: off Umhloti River; 73 m; 21 December 1900.  
Collected by: R.S. *Pieter Faure*.

#### Family Conidae

*Conus eucoronatus* Sowerby, 1903: 217, pl. 3 (fig. 9).

Holotype: SAM-A1753. Shell.

Locality: off Cape St. Blaize, 'N 85° W 4,5 miles'; 27 fm.; 15 June 1899.  
Collected by: R.S. *Pieter Faure*.

*Conus gilchristi* Sowerby 1903: 217, pl. 3 (fig. 8)

= *Conus natalis* Sowerby: Walls, 1979: 757.

Holotype: SAM-A1752. Shell.

Locality: Umhlangkulu River mouth, 'NW by N 7,5 miles'; 50 fm.; 14  
March 1901.

Collected by: R.S. *Pieter Faure*.

*Conus patens* Sowerby, 1903: 218, pl. 3 (fig. 7).

Holotype: SAM-A1756. Shell.

Locality: Vasco de Gama Peak, 'N 10° E 13 miles'; 85 fm.; 3 May 1900.  
Collected by: R.S. *Pieter Faure*.

#### Family Terebridae

*Terebra (Abretia) lightfooti* Smith, 1899: 247, pl. 6 (fig. 1).

Paratype: SAM-5519. Shell.

Locality: Table Bay; 10 fm.; no date.

Collected by: R. M. Lightfoot.

*Terebra planecosta* Barnard, 1958: 81, fig. 1a–b.

Holotype: SAM-A8659. Shell.

Locality: off Cove Rock, '4,5 miles'; 22 fm.; 6 August 1901.

Collected by: R.S. *Pieter Faure*.

Paratypes: SAM-A8658. Two shells.

Locality: off Cape St. Blaize, 'N by E 73 miles'; 125 fm.; 21 December 1899.

Collected by: R.S. *Pieter Faure*.

*Terebra thielei* Barnard, 1958: 80, fig. 1d.

Syntypes: SAM-A8657. Four shells.

Locality: Algoa Bay; 67 fm.; 10 November 1898.

Collected by: R.S. Pieter Faure.

#### Family Turridae

'*Acrobella*' *acus* Barnard, 1958: 149, fig. 21e.

Syntypes: SAM-A8748. Two shells, one fragment.

Locality: off Cape Recife, 34°27'S 25°42'E; 468 m; 16 November 1898.

Collected by: R.S. Pieter Faure.

'*Bela*' *bella* Barnard, 1958: 140, fig. 21c.

Syntypes: SAM-A8639. Three shells.

Locality: off Cape Agulhas; 146–183 m; 17 July 1903.

Collected by: R.S. Pieter Faure.

*Clavatula lobatopsis* Barnard, 1963c: 420, fig. 2a.

Syntypes: SAM-A9730. Three shells with opercula, radula slide.

Locality: 33°36'S 16°15'E; 2 780–2 880 m; August 1959.

Collected by: R.S. *Africana II*.

'*Cythara*' *dagama* Barnard 1963c: 423, fig. 2g–h.

Syntypes: SAM-A9731. Four shells, three with animals.

Locality: 33°36'S 16°15'E; 2 790–2 890 m; August 1959.

Collected by: R.S. *Africana II*.

'*Cythara*' *deliciosa* Barnard, 1958: 151, fig. 24

= *Eucithera deliciosa* (Barnard): Barnard, 1969: 610.

Syntypes: SAM-A8692. Six shells.

Locality: off Umhloti River; 73 m; 18 December 1900.

Collected by: R.S. Pieter Faure.

*Cythara*? *glaucocreas* Barnard, 1963c: 422, fig. 2d.

Holotype: SAM-A9821. Shell, radula slide.

Locality: 33°52'S 16°51'E; 2 430–2 790 m; December 1959.

Paratype: SAM-A9753. Shell.

Locality: 33°49'S 16°30'E; 2 750 m; August 1959.

Collected by: R.S. *Africana II*.

*Daphnella*? *bitrudis* Barnard, 1963c: 428, fig. 1b.

Syntypes: SAM-A9754. Two shells.

Locality: 33°49'S 16°30'E; 2 750 m; August 1959.

Collected by: R.S. *Africana II*.

*Daphnella recifensis* Barnard, 1958: 161, fig. 30a.

Holotype: SAM-A8757. Shell.

Locality: off Cape Recife, 34°27'S 25°42'E; 468 m; 14 November 1898.

Collected by: R.S. Pieter Faure.

*Daphnella? verecunda* Barnard, 1963c: 427, fig. 2e.

Syntypes: SAM-A9697. Three shells, one with animal.

Locality: 33°50'S 17°21'E; 1 100 m; August 1959.

Collected by: R.S. *Africana II*.

*Drillia armilla* Barnard, 1958: 137.

Holotype: SAM-A8714. Shell.

Locality: off Cove Rock; 146–239 m; 30 July 1901.

Collected by: R.S. Pieter Faure.

*Drillia bruchia* Barnard, 1958: 130, fig. 13b.

Holotype: SAM-A8717. Shell, apex missing.

Locality: off Cape Natal; 440 fm.; 4 April 1901.

Collected by: R.S. Pieter Faure.

*Drillia collina* Barnard, 1958: 129, fig. 13b.

Syntype: SAM-A8587. Shell.

Locality: off East London; 32 fm.; 1898.

Syntypes: SAM-A8710. Shell, two fragments.

Locality: off Hood Point; 49 fm.; 1901.

Collected by: R.S. Pieter Faure.

*Drillia diasi* Barnard, 1958: 137, fig. 19b.

Syntypes: SAM-A8566. Two shells.

Locality: 34°26'S 25°42'E; 124 fm.; 1898.

Collected by: R.S. Pieter Faure.

*Drillia dovyalis* Barnard, 1969: 606, fig. 6a.

Holotype: SAM-A29760. Shell.

Locality: off Sandy Point; 93 m; 14 August 1901.

Collected by: R.S. Pieter Faure.

*Drillia erepta* Barnard, 1969: 609, fig. b.

Syntype: SAM-A8734. Shell.

Locality: off Cape Morgan; 47 fm.; 25 July 1901.

Collected by: R.S. Pieter Faure.

Syntype: SAM-A29763. Shell.

Locality: 33°50'S 25°54'E (Algoa Bay); ?30 fm.; 11 November 1898.

Collected by: R.S. Pieter Faure.

Syntypes: SAM-A29764. Two shells.

Locality: off Cape St. Blaize; 125 fm.; 21 December 1899.  
Collected by: R.S. *Pieter Faure*.

*Drillia falcicosta* Barnard, 1958: 132, fig. 15b.

Syntypes: SAM-A8724. Two shells, one apex.  
Locality: off Umhloti River; 40 fm.; 1900.  
Collected by: R.S. *Pieter Faure*.

*Drillia laterculoides* Barnard, 1958: 129, fig. 12c.

Syntypes: SAM-A8709. Shell, apex.  
Locality: off Hood Point; 90 m; 15 July 1901.  
Collected by: R.S. *Pieter Faure*.

*Drillia latusulcus* Barnard, 1958: 131, fig. 13c.

Syntypes: SAM-A8718. Five shells.  
Locality: off Hood Point; 90 m; 15 July 1901.  
Collected by: R.S. *Pieter Faure*.

*Drillia morgana* Barnard, 1958: 133, fig. 16.

Holotype: SAM-A8739. Shell.  
Locality: off Cape Morgan; 47 fm.; 25 July 1901.  
Collected by: R.S. *Pieter Faure*.

*Drillia omia* Barnard, 1958: 123, fig. 11a.

Syntypes: SAM-A8651. Ten shells.  
Locality: ?Port Alfred, probably Table Bay-Saldanha Bay.  
Collector unknown.

*Drillia oneili* Barnard, 1958: 133, fig. 15c.

Holotype: SAM-A8731. Shell, slightly worn.  
Locality: off O'Neil Peak; 90 fm.; 1901.  
Collected by: R.S. *Pieter Faure*.

*Drillia pecus* Barnard, 1969: 607, fig. 5b.

Holotype: SAM-A29761. Shell.  
Locality: off East London; 400-450 fm.; 17 April 1901.  
Collected by: R.S. *Pieter Faure*.

*Drillia perfluans* Barnard, 1958: 132, fig. 15a.

Holotype: SAM-A8716. Shell.  
Locality: off Hood Point; 49 fm.; 15 July 1901.  
Collected by: R.S. *Pieter Faure*.

*Drillia pleonastica* Barnard, 1958: 133, fig. 17a.

Holotype: SAM-A8565. Shell.  
Locality: 34°26'S 25°42'E; 228 m; 14 November 1898.  
Collected by: R.S. *Pieter Faure*.

*Drillia pselia* Barnard, 1958: 138.

Syntypes: SAM-A8715. Shell, two broken shells.  
Locality: off Cape St. Blaize; 125 fm.; 1899.  
Collected by: R.S. Pieter Faure.

*Drillia simplicingula* Barnard, 1958: 135, fig. 18b.

Holotype: SAM-A8666. Shell.  
Locality: off Hood Point; 49 fm.; 15 July 1901.  
Paratype: SAM-A3471. Shell.  
Locality: off Cape Morgan; 104 m; 6 February 1901.  
Collected by: R.S. Pieter Faure.

*Drillia spiralis* Barnard, 1958: 134, fig. 17b.

Holotype: SAM-A8583. Shell.  
Locality: off Cape St. Blaize; 228 m; 21 December 1899.  
Collected by: R.S. Pieter Faure.

*Drillia tholos* Barnard, 1958: 130, fig. 14.

Syntypes: SAM-A8735. Fourteen shells.  
Locality: off Hood Point; 49 fm.; 15 July 1901.  
Collected by: R.S. Pieter Faure.

*Mangilia (Eucythara) africana* Sowerby, 1903: 21, pl. 5 (fig. 9)

= *Cythara africana* (Sowerby): Barnard, 1958: 115.

Holotype: SAM-A345. Shell.  
Paratype: SAM-A33954. Shell, broken, outer lip thin.  
Locality: off Umhloti River; 46 m; 18 December 1900.  
Collected by: R.S. Pieter Faure.

*Mangilia extans* Barnard, 1958: 156, fig. 27c.

Holotype: SAM-A8756. Shell.  
Locality: off Cove Rock; 40 m; 6 August 1901.  
Collected by: R.S. Pieter Faure.

*Mangilia muiri* Barnard, 1958: 153, fig. 26a.

Syntypes: SAM-A8647. Seven shells, in growth series.  
Locality: Still Bay; intertidal; 1921.  
Muir Collection.

*Mangilia phoxos* Barnard, 1958: 155, fig. 27b.

Holotype: SAM-A8730. Shell.  
Locality: off Umhloti River; 73 m; 18 December 1900.  
Collected by: R.S. Pieter Faure.



*Mangilia sciola* Barnard, 1958: 157, fig. 28c.

Syntype: SAM-A8642. Shell.

Locality: off Cape Recife, 34°27'S 25°42'E; 468 m; 14 November 1898.

Syntype: SAM-A8643. Shell.

Locality: near East London, 33°06'S 27°55'E; 78 m; 14 November 1898.

Collected by: R.S. Pieter Faure.

*Mangilia serrula* Barnard, 1964c: 15, fig. 1a-b.

Syntypes: SAM-A8728. Five shells.

Locality: off Illovo and Umkomaas; 30-40 fm.; 31 December 1900.

Collected by: R.S. Pieter Faure.

*Mangilia tranquilla* Barnard, 1958: 158, fig. 28b.

Syntypes: SAM-A8648. Five shells.

Locality: Still Bay; intertidal; no date.

Collected by: J. Muir.

*Mangilia translucens* Barnard, 1958: 155, fig. 27a.

Syntypes: SAM-A8586. Twelve shells.

Locality: off Umhloti River mouth; 40 fm.; 28 December 1898.

Collected by: R.S. Pieter Faure.

*Mitramorpha veneris* Barnard, 1964c: 16.

Holotype: SAM-A8750. Shell, broken.

Locality: off Cape Vidal; 146-183 m; 27 February 1901.

Collected by: R.S. Pieter Faure.

*Monilopsis psilarosis* Barnard, 1963c: 421, fig. 4a.

Syntypes: SAM-A9789. Two shells, radula slide.

Locality: 34°42'S 16°54'E; 3 140-3 270 m; December 1959.

Collected by: R.S. *Africana II*.

*Philbertia natalensis* Barnard, 1958: 158, fig. 29a.

Syntype: SAM-A8654. Shell.

Locality: off Umhloti River mouth; 40 fm.; 18 December 1900.

Collected by: R.S. Pieter Faure.

Syntype: SAM-A8758. Shell.

Locality: off Cape Natal, 54 fm.; 14 December 1900.

Collected by: R.S. Pieter Faure.

*Pleurotoma (Genotia) belaeformis* Sowerby, 1903: 216, pl. 4 (fig. 8)

= '*Genotia*' *belaeformis* (Sowerby): Barnard, 1958: 112, figs 8b-9b.

Paratypes: SAM-A1687. Two shells.

Locality: off Cape Point; 230 fm.; 4 May 1900.

Collected by: R.S. Pieter Faure.

'*Pleurotoma*' *curricula* Barnard, 1969: 612, fig. 8.

Syntypes: SAM-A9859. Shell, two broken.

Locality: 34°36'S 17°00'E; 2 750-3 200 m; December 1959.

Collected by: R.S. *Africana II*.

*Pleurotoma* (*Drillia*) *fossata* Sowerby, 1903: 214, pl. 3 (fig. 5)

= *Drillia fossata* (Sowerby): Barnard, 1958: 135.

Holotype: SAM-A346. Shell.

Paratype: SAM-A346. Shell.

Locality: off Cape Vidal; 80-100 fm.; 27 February 1901.

Collected by: R.S. *Pieter Faure*.

*Pleurotoma* (*Clavus*) *lignaria* Sowerby, 1903: 215, pl. 3 (fig. 4)

= *Turris lignaria* (Sowerby): Barnard, 1958: 102.

Paratype: SAM-A357. Shell.

Locality: off Table Bay; 240 m; 28 March 1900.

Collected by: R.S. *Pieter Faure*.

*Pleurotoma* (*Surcula*) *lobata* Sowerby, 1903: 213, pl. 4 (fig. 9)

= *Turris lobata* (Sowerby): Barnard, 1958: 107, fig. 3i-j.

Syntypes: SAM-A1673. Five shells, radula slide.

Locality: off Cape Natal; 810 m; 4 April 1901.

Syntypes: SAM-A1674. Three shells, one with operculum.

Locality: off Buffalo River, East London; 560 m; 24 April 1901.

Collected by: R.S. *Pieter Faure*.

*Pleurotoma* (*Drillia*) *scitecostata* Sowerby, 1903: 214, pl. 4 (fig. 10)

= '*Drillia*' *scitecostata* (Sowerby): Barnard, 1958: 119, figs 8f, 20.

Syntypes: SAM-A347. Two shells.

Locality: off Glendower Beacon; 100 fm.; 10 September 1901.

Collected by: R.S. *Pieter Faure*.

*Pleurotoma* (*Clavatula*) *turriplana* Sowerby, 1903: 215, pl. 3 (fig. 6)

= *Clavatula turriplana* Sowerby: Barnard, 1958: 144.

Holotype: SAM-A1646. Shell.

Locality: off Cape St. Blaize, 'N by E  $\frac{1}{4}$  E 65 miles'; 155 m; 22 December 1899.

Collected by: R.S. *Pieter Faure*.

*Surcula aditus* Barnard, 1969: 604, fig. 4a.

Holotype: SAM-A29759. Shell.

Locality: off Cove Rock; 22 fm.; 6 August 1901.

Collected by: R.S. *Pieter Faure*.

*Surcula amplisulcus* Barnard, 1958: 147, fig. 22a.

Syntypes: SAM-A1689. Five shells.

Locality: west coast of Cape Peninsula; 120 fm.; 4 April 1902.

Collected by: R.S. Pieter Faure.

*Surcula faurei* Barnard, 1958: 147, fig. 22b.

Holotype: SAM-A8611. Shell.

Locality: Brown's Bank, 36,5°S 21°E; 80–100 fm.; 17 July 1906.

Collected by: R.S. Pieter Faure.

*Surcula scalaria* Barnard, 1958: 146, fig. 22d.

Syntypes: SAM-A358. Seven shells.

Locality: off Cape Point; 480–800 fm.; 10 September 1903.

Collected by: R.S. Pieter Faure.

*Surcula sulcancellata* Barnard, 1958: 145, fig. 22c.

Syntypes: SAM-A361. Five shells.

Locality: off Cape Point; 130–300 fm.; 28 September 1903.

Collected by: R.S. Pieter Faure.

*Turris ambages* Barnard, 1958: 148, fig. 23b.

Syntypes: SAM-A8683. Two shells.

Locality: off Cape Natal; 54 fm.; 14 December 1900.

Collected by: R.S. Pieter Faure.

Syntype: SAM-A8684. Shell.

Locality: off Umkomaas River; 40 fm.; 31 December 1900.

Collected by: R.S. Pieter Faure.

Syntype: SAM-A8685. Shell.

Locality: off O'Neil Peak; 90 fm.; 28 February 1901.

Collected by: R.S. Pieter Faure.

*Turris saldanhae* Barnard, 1958: 109, figs 3f, 7.

Syntypes: SAM-A1738. Seventeen shells.

Locality: Baboon Point, Saldanha Bay; 31 fm.; 18 March 1902.

Collected by: R.S. Pieter Faure.

*Typhlomangilia? polythele* Barnard, 1963c: 422, fig. 3e–f.

Syntypes: SAM-A9802. Two shells.

Locality: 33°50'S 16°30'E; 1 480–1 660 fm.; December 1950.

Collected by: F. H. Talbot.

*Typhlosyrinx chrysoplex* Barnard, 1963c: 425, fig. 3g.

Holotype: SAM-A9857. Shell.

Locality: 34°36'S 17°00'E; 1 500–1 760 fm.; December 1959.

Collected by: F. H. Talbot.

*Typhlosyrinx pyrropelex* Barnard, 1963c: 423, fig. 2c.

Syntypes: SAM-A1643. Two shells.

Locality: off Cape Point, N 77° E; 650–700 fm.; 29 July 1903.

Collected by: R. S. Pieter Faure.

Syntypes: SAM-A9858. Three shells, juveniles with animals.

Locality: 34°36'S 17°00'E; 1 500–1 760 fm.; December 1959.

Collected by: F. H. Talbot.

*Typhlosyrinx subrosea* Barnard, 1963c: 425, fig. 3a–d.

Holotype: SAM-A9822. Shell with animal.

Locality: 33°52'S 16°51'E; 1 380–1 520 fm.; December 1959.

Collected by: F. H. Talbot.

Paratype: SAM-A9822. Shell.

Locality: 33°52'S 16°51'E; 1 380–1 520 fm.; December 1959.

Collected by: F. H. Talbot.

#### Subclass OPISTHOBRANCHIA

##### Family Acteonidae

*Solidula niecaensis* Barnard, 1963d: 317

= *Pupa niecaensis* (Barnard): **comb. nov.**

Holotype: SAM-A6553. Shell.

Locality: off Nieca River; 80 m; 7 August 1901.

Collected by: R.S. Pieter Faure.

##### Family Retusidae

*Retusa natalensis* Barnard, 1963d: 321, fig. 28a.

Syntypes: SAM-A9409. Two shells.

Locality: off Umkomaas River; 40 fm.; 1900.

Collected by: R.S. Pieter Faure.

*Volvula mutabilis* Barnard, 1963d: 322, fig. 28b.

Syntypes: SAM-A9410. Two shells.

Locality: off Umhloti River; 40 fm.; 1900.

Collected by: R.S. Pieter Faure.

##### Family Scaphandridae

*Acteocina natalensis* Barnard, 1963d: 324, fig. 28c.

Syntypes: SAM-A9388. Two shells.

Locality: off Illovo River; 27–30 fm.; 1900.

Collected by: R.S. Pieter Faure.

Family **Doridoxidae**

*Doridoxa benthalis* Barnard, 1963c: 443, fig. 10.

Holotype: SAM-A9796. Animal, preserved, jaws, radula slide.

Locality: 34°42'S 16°54'E; 1 725–1 780 fm.; December 1959.

Collected by: F. H. Talbot.

Family **Chromodorididae**

*Glossodoris capensis* Barnard, 1927: 181, pl. 19 (figs 3–4)

= *Hypselodoris capensis* (Barnard): **comb. nov.**

Syntypes: SAM-A6230. Two animals, preserved, two pairs jaws, two radula slides.

Locality: St. James, False Bay; intertidal; April 1912.

Collected by: K. H. Barnard.

Family **Polyceridae**

*Polycera nigrocrocea* Barnard, 1927: 191, pl. 19 (figs 7–8)

= *Polycera capensis* Quoy & Gaimard: Pruvot-Fol, 1934: 75.

Syntype: SAM-A6225. Two animals, preserved, radula.

Locality: St. James; intertidal; March 1912.

Collected by: K. H. Barnard.

*Plocamopherus apheles* Barnard, 1927: 195, fig. 1.

Holotype: SAM-A6232. Animal, preserved, jaws, radula slide.

Locality: Nahoon estuary, East London; no depth; 31 May 1901.

Collected by: J. D. F. Gilchrist.

Family **Goniodorididae**

*Goniodoris ovata* Barnard, 1934: 2, 2 text figures.

Holotype: SAM-A6270. Animal, preserved.

Paratype: SAM-A6270. Radula slide.

Locality: Simonstown; from boat hull; June 1933.

Collected by: K. H. Barnard.

Family **Janolidae**

*Bonisa nakaza* Gosliner, 1981: 19, figs 1d–f, 18–25.

Holotype: SAM-A34886. Animal, preserved.

Paratypes: SAM-A34887. Four animals, preserved.

Locality: off Llandudno, 34°01'S 18°20'E; 20 m; 23 December 1979.

Collected by: W. R. Liltved.

*Janolus longidentatus* Gosliner, 1981: 9, figs 1c, 10–17.

Holotype: SAM-A34883. Animal, preserved.

Locality: Miller's Point, 34°14'S 18°29'E; 3 m; 10 December 1980.

Collected by: T. M. Gosliner.

Paratypes: SAM-A34884. Two animals, preserved.

Locality: Castle Rocks, 34°18'S 18°29'E; 10 m; 1 October 1980.

Collected by: T. M. Gosliner.

Paratype: SAM-A34885. Animal, preserved.

Locality: Miller's Point, 34°14'S 18°29'E; 5 m; 11 July 1980.

Collected by: T. M. Gosliner.

#### Family **Flabellinidae**

*Flabellina funeka* Gosliner & Griffiths, 1981: 106, figs 1a, 2–4.

Holotype: SAM-A34317. Animal, preserved.

Locality: Castle Rocks, 34°14'S 18°29'E; 10 m; 17 January 1980.

Collected by: T. M. Gosliner.

Paratype: SAM-A34318. Animal, preserved.

Locality: Castle Rocks, 34°14'S 18°29'E; 10 m; 17 January 1980.

Collected by: T. M. Gosliner.

Paratype: SAM-A34319. Animal, preserved.

Locality: Venus Pool, 34°17'S 18°28'E; 10 m; 6 January 1980.

Collected by: T. M. Gosliner.

Paratype: SAM-A34320. Animal, preserved.

Locality: Hermanus, 34°17'S 19°15'E; 17 m; 11 October 1971.

Collected by: R. J. Griffiths.

Paratype: SAM-A34321. Animal, preserved.

Locality: Hermanus, 34°17'S 19°15'E; 17 m; 11 October 1971.

Collected by: R. J. Griffiths.

#### Family **Tergipedidae**

*Catriona casha* Gosliner & Griffiths, 1981: 130, figs 12–14.

Holotype: SAM-A34871. Animal, preserved.

Paratypes: SAM-A34872. Five animals, preserved.

Locality: Cape Town docks, 33°54'S 18°26'E; 1 m; 26 June 1972.

Collected by: R. J. Griffiths.

#### Family **Facelinidae**

*Facelina faurei* Barnard, 1927: 205, figs. 4–5

= *Pruvotfolia faurei* (Barnard): Tardy, 1969: 343.

Syntypes: SAM-A6249. Several animals, preserved, jaws, radula slide.

Locality: off Saldanha Bay; 33 fm.; 12 March 1902.

Collected by: R.S. Pieter Faure.

*Favorinus horridus* Macnae, 1954: 19, pl. 1 (fig. 4) figs 11–13

= *Phyllodesmium serratum* (Baba): Baba, 1955: 56.

Paratypes: SAM–A29578. Fifteen animals, preserved.

Locality: '½ mile S of Kalk Bay Harbour'; intertidal; 1949.

Collected by: W. Macnae.

*Hervia quadricolor* Barnard, 1927: 203, pl. 20 (figs 9–10)

= *Godiva quadricolor* (Barnard): Macnae, 1954: 23.

Holotype: SAM–A6248. Animal, preserved.

Locality: St. James (False Bay); intertidal; 20 June 1912.

Collected by: K. H. Barnard.

#### Family Cratenidae

*Cratena capensis* Barnard, 1927: 204, pl. 20 (fig. 8).

Syntype: SAM–A6247. Jaws, radula slide.

Locality: Sea Point; intertidal; 29 November 1913.

Collected by: K. H. Barnard.

#### Family Aeolidiidae

*Aeolidiella saldanhensis* Barnard, 1927: 201, figs 2–3

= *Aeolidiella indica* Bergh: Gosliner & Griffiths, 1981: 119.

Syntype: SAM–A6257. Jaws, radula slide.

Locality: Saldanha Bay; intertidal; 5 September 1912.

Collected by: R.S. Pieter Faure.

#### Subclass PULMONATA

#### Family Siphonariidae

*Siphonaria (Patellopsis) anneae* Tomlin, 1944: 92, text figure.

Syntypes: SAM–A2988. Two shells.

Locality: Umhlali; no depth or date.

Collected by: H. C. Burnup.

*Siphonaria cyaneomaculata* Sowerby, 1906: 37, text figure

= *Siphonaria (Patellopsis) deflexa* (Helbling): Allanson, 1958: 159.

Syntypes: SAM–A3330. Two shells.

Locality: Kowie, Port Alfred; intertidal; 1915.

Collected by: H. Becker.

*Siphonaria (Patellopsis) dayi* Allanson, 1958: 169, pl. 16 (figs 10–15), text figure.

Holotype: SAM–A32707. Shell.

Locality: Delagoa Bay; no depth or date.

Collected by: University of Cape Town.

Paratypes: SAM–A33403. Three shells.

Locality: Inhaca Island; no depth; 30 December 1958.

Collected by: University of Cape Town.

*Siphonaria (Siphonaria) aspera pallida* Allanson, 1958: 171, pl. 4, fig. 17.

Syntypes: SAM–A33402. Four shells.

Locality: Langebaan Lagoon; intertidal; 25 April 1949.

Collected by: University of Cape Town.

#### Class SCHAPHOPODA

#### Family Dentaliidae

*Dentalium africanum* Sowerby, 1903: 224, pl. 5 (fig. 10).

Paratypes: SAM–A5489. Six shells, radula slide.

Locality: off Umtwalumi River; 25 fm.; 22 March 1901.

Collected by: R.S. Pieter Faure.

*Dentalium (Compressidens) capense* Tomlin, 1931: 340.

Holotype: SAM–A6191. Shell.

Locality: off Cape Point; 900 fm.; 19 August 1903.

Collected by: R.S. Pieter Faure.

*Dentalium engischistum* Barnard, 1963d: 352, fig. 30f.

Syntypes: SAM–A5463. Six shells, two with animals, one broken, radula slide.

Locality: off Cape Natal; 62 fm.; 1900.

Collected by: R.S. Pieter Faure.

*Dentalium eualdes* Barnard, 1963c: 444.

Syntypes: SAM–A9736. Two shells, radula slide.

Locality: 33°36'S 16°15'E; 2 780–2 880 m; August 1959.

Collected by: R.S. *Africana II*.

*Dentalium lardum* Barnard, 1963c: 445.

Syntypes: SAM–A9768. Three shells.

Locality: 33°39'S 16°30'E; 1 500 fm.; August 1959.

Collected by: R.S. *Africana II*.



*Dentalium natalense* Barnard, 1963d: 350, fig. 30e.

Syntypes: SAM-A9364. Forty-nine shells.

Locality: off Cape Natal; 85 fm.; 17 December 1900.

Collected by: R.S. Pieter Faure.

*Dentalium (Fissidentalium) platypleurum* Tomlin, 1931: 339, text figure.

Holotype: SAM-A3631. Shell.

Locality: off Itongazi River; 25 fm.; 14 March 1901.

Collected by: R.S. Pieter Faure.

*Dentalium salpinx* Tomlin, 1931: 338, text figure.

Holotype: SAM-A5459. Shell.

Paratypes: SAM-A5459. Nine shells.

Locality: off Cape Point, 'NE  $\frac{3}{4}$  E, 40 miles'; 700-800 fm.; 15 September 1903.

Collected by: R.S. Pieter Faure.

*Dentalium subterlineatum* Tomlin, 1931: 337.

Holotype: SAM-A6192. Shell.

Locality: off Cape Point; 900 fm.; 19 August 1903.

Collected by: R.S. Pieter Faure.

#### Family Cadulidae

*Cadulus promontorii* Barnard, 1963d: 353, fig. 30h-l.

Syntypes: SAM-A7460. Four shells, radula slide.

Locality: off Cape Point; 700 fm.; 20 August 1903.

Collected by: R.S. Pieter Faure.

#### Class BIVALVIA

#### Family Nuculidae

*Nucula aequalitas* Barnard, 1964a: 365.

Holotype: SAM-A9472. Two valves.

Locality: off East London; 732-915 m; 17 April 1901.

Collected by: R.S. Pieter Faure.

*Nucula fragilis* Boshoff, 1968: 95, pl. 8a, text fig. 1.

Syntypes: SAM-A30212. Two valves.

Locality: 29°34'S 31°39'E; 115-118 m; 9 September 1964.

Collected by: R.V. Anton Bruun.

*Nucula irregularis* Sowerby, 1904: 7, pl. 6 (fig. 12).

Paratypes: SAM-14843. Two valves.

Locality: off Struis Point; 28 m; 17 July 1902.

Collected by: R.S. *Pieter Faure*.

*Nucula sculpturata* Sowerby, 1904: 7, pl. 6 (fig. 11)

= *Nucula pulchra* Hinds: Barnard, 1964a: 362.

Paratypes: SAM-14830. Ten valves, three with dry animals.

Locality: 33°03'S 27°57'E; 62 m; 28 December 1898.

Collected by: R.S. *Pieter Faure*.

#### Family Mallettiidae

*Malletia estheriopsis* Barnard, 1963c: 447, fig. 11b.

Syntypes: SAM-A9817. Four valves.

Locality: 33°50'S 16°30'E; 2 700–3 040 m; December 1959.

Collected by: F. H. Talbot on R.S. *Africana II*.

#### Family Nuculanidae

*Leda lanceta* Boshoff, 1968: 96, pl. 8b–c.

Holotype: SAM-A30213. Two valves.

Locality: 29°21'S 31°58'E; 370 m; no date.

Collected by: R.V. *Anton Bruun*.

*Leda macella* Barnard, 1963c: 448, fig. 11d

= *Nuculana vestita* (Locard): Knudsen, 1970: 28, fig. 11.

Syntypes: SAM-A9851. Sixty-six complete shells, nine valves.

Locality: 34°05'S 16°58'E; 2 690–2 720 m; December 1959.

Collected by: F. H. Talbot on R.S. *Africana II*.

*Leda parceplicata* Barnard, 1964b: 21

= *Portlandia (Portlandia) parceplicata* (Barnard): Kilburn, 1973b: 698.

Syntypes: SAM-A9465. Thirty-nine complete shells, 157 valves.

Locality: off Umhloti River, Natal; 73 m; 18 December 1900.

Collected by: R.S. *Pieter Faure*.

*Leda parsimonia* Barnard, 1963c: 448.

Holotype: SAM-A9818. Two valves.

Locality: 33°50'S 16°30'E; 2 700–3 040 m; August 1959.

Collected by: F. H. Talbot on R.S. *Africana II*.

*Nuculana compta* Sowerby, 1904: 6, pl. 6 (fig. 10)

= *Leda compta* (Sowerby): Barnard, 1964a: 367, fig. 1a.

Paratype: SAM-14818. Two valves.

Locality: off Cape Natal; 805 m; 4 April 1901.

Collected by: R.S. *Pieter Faure*.

*Nuculana gemmulata* Sowerby, 1904: 6, pl. 6 (fig. 9)

= *Leda gemmulata* (Sowerby): Barnard 1964a: 366.

Paratypes: SAM-14785. Four valves.

Locality: off Tugela River; 68 m; 29 January 1901.

Collected by: R.S. Pieter Faure.

*Nuculana lamellata* Sowerby, 1904: 5, pl. 6 (fig. 8)

= *Leda lamellata* (Sowerby): Barnard, 1964a: 366.

Paratypes: SAM-14787. Sixteen complete shells, thirteen valves.

Locality: off Cape Natal; 99 m; 14 December 1900.

Collected by: R.S. Pieter Faure.

*Sarepta natalensis* Barnard, 1964c: 22, fig. 4a.

Holotype: SAM-A9474. Valve.

Locality: off Cape Natal; 90 m; 14 December 1900.

Paratypes: SAM-A9470. Seven valves.

Locality: off Tugela River, Natal; 110-146 m; 11 January 1901.

Collected by: R.S. Pieter Faure.

#### Family Arcidae

*Arca (Scapharca) africana* Sowerby, 1904: 4, pl. 6 (fig. 4).

Paratypes: SAM-14831. Complete shell with animal, two valves.

Locality: off Tugela River; 84-100 m; 6 February 1901.

Collected by: R.S. Pieter Faure.

*Deltaodon tugelae* Barnard, 1962b: 249, figs 2a-e. Kilburn, 1973b: 699.

Lectotype: SAM-A31736. One valve.

Paralectotypes: SAM-A9477. Eighteen valves.

Locality: off Zululand; no depth or date.

Collected by: R.S. Pieter Faure.

#### Family Limopsidae

*Limopsis natalis* Barnard, 1964b: 23, figs.

Syntypes: SAM-A9479. Complete shell, two valves.

Locality: off O'Neil Peak, Natal; 103 m; 28 February 1901.

Collected by: R.S. Pieter Faure.

#### Family Mytilidae

*Crenella striatissima* Sowerby, 1904: 3, pl. 6 (fig. 1). Barnard 1964a: 403.

Holotype: SAM-14855. Two valves.

Locality: 37°07'S 25°40'E; 106 m; 14 November 1898.

Paratype: SAM-14856. Valve, broken.

Locality: off Umhloti River, Natal; 183 m; 19 December 1900.  
Collected by: R.S. Pieter Faure.

*Gregariella simplicifilis* Barnard, 1964a: 402.

Syntypes: SAM-6878. Five complete shells, animals in three, five valves.  
Locality: Gordon's Bay, False Bay; intertidal; no date.  
Collected by: R. M. Lightfoot.

*Musculus (Musculus) virgiliae* Barnard, 1964a: 399, fig. 8d.

Syntypes: SAM-A7776. Six complete shells, two valves, one broken.  
Locality: Keurbooms River, eastern Cape; intertidal; 1931.  
Collected by: K. H. Barnard.

*Septifer bisculpturata* Barnard, 1964c: 23, fig. 4c-d.

Holotype: SAM-A9482. Valve.  
Locality: off Tugela River, Natal; 86 m; 29 January 1901.  
Collected by: R.S. Pieter Faure.

#### Family Pectinidae

*Chlamys fultoni* Sowerby, 1904: 2, pl. 6 (fig. 5).

Paratype: SAM-14970. Valve.  
Locality: off Amatikulu River, Natal; 48 m; 1 February 1901.  
Collected by: R.S. Pieter Faure.

*Chlamys gilchristi* Sowerby, 1904: 1, pl. 6 (fig. 6). Barnard 1964a: 427.

Holotype: SAM-14853. Shell, complete.  
Locality: False Bay; 416 m; 4 May 1900.  
Collected by: R.S. Pieter Faure.

*Chlamys humilis* Sowerby, 1904: 3, pl. 6 (fig. 3)

= *Chlamys tinctus* (Reeve): Barnard, 1964a: 425.

Paratype: SAM-14858. Shell, complete.  
Locality: off Cape St. Blaize, Mossel Bay; 167-183 m; 20 December 1900.  
Collected by: R.S. Pieter Faure.

*Cyclopecten incubans* Barnard, 1964a: 432.

Syntypes: SAM-A9493. Twelve valves.  
Locality: off Cape St. Blaize; 230 m; 21 December 1899.  
Collected by: R.S. Pieter Faure.

*Cyclopecten vidalensis* Barnard, 1964a: 433, fig. 14e-g.

Holotype: SAM-A9495. Valve.  
Locality: off Cape Vidal, Zululand; 146-183 m; 27 February 1901.  
Collected by: R.S. Pieter Faure.

Family **Limidae**

*Lima abscisa* Barnard, 1964a: 441, fig. 16f.

Syntypes: SAM-A9504. Three valves.

Locality: off Cape Morgan, near East London; 86 m; 25 July 1901.

Collected by: R.S. Pieter Faure.

*Lima divericata* Barnard, 1964a: 440, fig. 16e.

Syntypes: SAM-A9501. Two valves.

Locality: off O'Neil Peak, Zululand; 165 m; 28 February 1901.

Collected by: R.S. Pieter Faure.

*Lima symmetrica* Barnard, 1964a: 441.

Holotype: SAM-A9503. Valve.

Locality: off Cape Vidal; 146-183 m; 27 February 1901.

Collected by: R.S. Pieter Faure.

Family **Astartidae**

*Astarte (Digitaria) lunulata* Barnard, 1964a: 449, fig. 18b.

Syntypes: SAM-A9506. Three valves.

Locality: 34°26'S 25°42'E; 226 m; 14 November 1898.

Collected by: R.S. Pieter Faure.

Family **Crassatellidae**

*Crassatella natalensis* Barnard, 1964a: 457, fig. 19b.

Syntypes: SAM-A9509. Four valves, one juvenile.

Locality: off Cape Natal; 98 m; 14 December 1900.

Collected by: R.S. Pieter Faure.

*Crassatella pilula* Barnard, 1964a: 457, fig. 19c.

Syntypes: SAM-A3665. Three complete shells, five valves.

Locality: off Cape Morgan, N of East London; 144 m; 26 July 1901.

Collected by: R.S. Pieter Faure.

Family **Carditidae**

*Cardita pulcherrima* Sowerby, 1904: 7, pl. 6 (fig. 2). Barnard, 1964a: 459.

Paratypes: SAM-A9512. Nine valves.

Locality: off Cape Natal; 98 m; 14 December 1900.

Collected by: R.S. Pieter Faure.

*Venericardia fortisculpta* Barnard, 1964a: 462.

Syntypes: SAM-A9519. Four valves.

Locality: Algoa Bay; 95 m; 14 November 1898.

Collected by: R.S. Pieter Faure.

*Venericardia nuculoides* Barnard, 1964a: 462.

Syntypes: SAM-A29661. Twenty-two complete shells, eight valves.

Locality: 35°05'S 18°17'E; 27 m; 2 July 1961.

Collected by: University of Cape Town.

#### Family Lucinidae

*Phacoides peritaphros* Barnard, 1964a: 476, fig. 22.

Syntypes: SAM-A9524. Five valves.

Locality: off Nieca River, East London; 43 fm.; 7 August 1901.

Collected by: R.S. Pieter Faure.

*Phacoides saldanhae* Barnard, 1964a: 474, fig. 21c.

Syntypes: SAM-A4473. Four valves.

Locality: off Baboon Point, Saldanha Bay; 56 m; 18 March 1902.

Collected by: R.S. Pieter Faure.

*Phacoides sepes* Barnard, 1964c: 25.

Syntypes: SAM-A9529. Two valves.

Locality: off Morewood Cove, Natal; 49 m; 19 December 1900.

Collected by: R.S. Pieter Faure.

*Phacoides sudes* Barnard, 1964c: 25, fig. 6a-b

= *Gonimyrtea sudes* (Barnard): Kilburn, 1973b: 701.

Syntypes: SAM-A9531. Six valves.

Locality: off Cape Natal; 98 m; 14 December 1900.

Collected by: R.S. Pieter Faure.

#### Family Laternulidae

*Periploma fractura* Boshoff, 1968: 97, pl. 8d, text-fig. 2.

Holotype: SAM-A30211. Shell, complete.

Locality: 29°29'S 30°41'E; 86 m; 9 September 1969.

Collected by: R.V. Anton Bruun.

#### Family Thyasiridae

*Thyasira unilateralis* Barnard, 1964c: 24, fig. 5a.

Syntypes: SAM-A9520. Two valves.

Locality: off O'Neil Peak, Natal; 165 m; 28 February 1901.

Collected by: R.S. Pieter Faure.

Family **Erycinidae**

*Tellimya biradialis* Barnard, 1964a: 484, fig. 23a.

Syntypes: SAM-A9535. Two valves.

Locality: off Cape Natal; 155 m; 17 December 1900.

Collected by: R.S. Pieter Faure.

*Tellimya trigona* Barnard, 1964a: 484, fig. 23b.

Syntypes: SAM-A29741. Five valves.

Locality: Langebaan (Saldanha Bay); no depth; 26 April 1949.

Collected by: University of Cape Town.

Family **Montacutidae**

*Conchentopyx granulosa* Barnard, 1964b: 35, fig. 2.

?Holotype: SAM-A33071. Animal, preserved.

Locality: Langebaan (Saldanha Bay); no depth or date.

Collected by: University of Cape Town.

*Montacula ornata* Barnard, 1964b: 26, fig. 6c-d

= *Barrimysia* (*Callomysia*) *ornata* (Barnard): Kilburn, 1973b: 702.

Syntypes: SAM-A9538. Three valves.

Locality: off Umhloti River; 73 m; 18 December 1900.

Collected by: R.S. Pieter Faure.

*Montacuta siliqua* Barnard, 1964b: 26, fig. 6e

= *Nippon mysella* (Barnard): Kilburn, 1973b: 702.

Syntypes: SAM-A9539. Three valves.

Locality: off Umhloti River; 73 m; 18 December 1900.

Collected by: R.S. Pieter Faure.

Family **Veneridae**

*Venus* (*Anaitis*) *intersculpta* Sowerby, 1904: 11, pl. 7 (fig. 2)

= *Venus verrucosa* Linnaeus: Barnard, 1964a: 496.

Paratype: SAM-14841. Shell, complete.

Locality: Algoa Bay; 18-29 m; 15 March 1899.

Collected by: R.S. Pieter Faure.

Family **Mesodesmatidae**

*Donacilla delagoae* Barnard, 1964a: 515, fig. 28b.

Syntypes: SAM-A29742. Two complete shells, juvenile, ten valves.

Locality: Inhaca Island, Delagoa Bay; no depth or date.

Collected by: University of the Witwatersrand.

Family **Tellinidae**

*Tellina acropisthus* Barnard, 1964b: 26. fig. 5b

= *Tellina (Tellinella) staurella* Lamarck: Boss, 1969: 95, pl. 2 (fig. 2), pl. 3 (figs 1-2), pl. 4 (fig. 2).

Syntype: SAM-A9547. Valve.

Locality: off Cape Natal; 88 m; 17 December 1900.

Collected by: R.S. Pieter Faure.

*Tellina europisthus* Barnard, 1964b: 27, fig. c

= *Tellina (Cadella) semen* Hanley: Boss, 1969: 136.

Syntypes: SAM-A9549. Four valves.

Locality: off Cape Natal; 98 m; 14 December 1900.

Collected by: R.S. Pieter Faure.

*Tellina gilchristi* Sowerby, 1904: 12, pl. 7 (fig. 3). Barnard, 1964a: 540

= *Tellina (Moerella) gilchristi* Sowerby: Boss, 1969: 144.

Paratypes: SAM-14751. Six valves.

Locality: off Cape Point; 92 m; 6 June 1900.

Collected by: R.S. Pieter Faure.

*Tellina (Macoma) inclinata* Sowerby, 1904: 14, pl. 7 (fig. 9)

= *Macoma inclinata* (Sowerby): Barnard 1964a: 549, fig. 31i.

Syntypes: SAM-14835. Six valves.

Locality: off Tugela River; 84-92 m; 6 February 1901.

Collected by: R.S. Pieter Faure.

*Tellina (Macoma) levior* Sowerby, 1904: 13, pl. 7 (fig. 6)

= *Macoma levior* (Sowerby): Barnard, 1964a: 548.

Paratype: SAM-14797. Four complete shells, 8 valves.

Locality: Tugela River, 'N by W 4 miles'; 24 fm.; 21 January 1901.

Collected by: R.S. Pieter Faure.

Paratype: SAM-14798. Six complete shells, 2 valves.

Locality: Amatakulu River, NW by W  $\frac{3}{4}$  W 12 miles; 26 fm.; 7 February 1901.

Collected by: R.S. Pieter Faure.

*Tellina (Macoma) ordinaria* Sowerby, 1904: 14, pl. 7 (fig. 7)

= *Macoma ordinaria* (Sowerby): Barnard, 1964a: 547, fig. 31e.

Paratypes: SAM-14824. Three valves.

Locality: off Saldanha Bay; 18-26 m; 19 March 1902.

Collected by: R.S. Pieter Faure.

*Tellina vidalensis* Sowerby, 1904: 13, pl. 7 (fig. 6). Barnard 1964a: 541

= *Tellina (Moerella) vidalensis* Sowerby: Boss, 1969: 141, pl. 16 (figs 2-5).



Holotype: SAM-14848. Shell, complete.

Locality: off Cape Vidal; 23 m; no date.

Collected by: R.S. Pieter Faure.

#### Family Pandoridae

*Pandora dissimilis* Sowerby, 1894: 374; 1897: 21, pl. 6 (fig. 33). Barnard, 1964a: 372.

Paratypes: SAM-5622. Two shells, complete.

Locality: Table Bay, Green Point; intertidal; no date.

Collected by: E. L. Layard.

*Pandora similis* Sowerby, 1897: 29. Barnard, 1964a: 572.

Paratypes: SAM-A29743. Two shells, complete.

Locality: Bluff Channel, Durban; no depth or date.

Donated by: Mr Ponsonby.

#### Family Cuspidariidae

*Cuspidaria nasuta* Sowerby, 1904: 18, pl. 7, fig. 14

= *Cuspidaria capensis* (Smith): Barnard, 1964a: 580.

Holotype: SAM-14819. Shell, complete.

Paratype: SAM-A4432. Shell, complete.

Locality: off Cape Point Lighthouse; 155 m; 3 May 1900.

Collected by: R.S. Pieter Faure.

*Cuspidaria optima* Sowerby, 1904: 17, pl. 11 (fig. 16). Barnard, 1964a: 580.

Paratype: SAM-14764. Shell, complete.

Locality: off Umtwalumi River, Natal; 92 m; 11 March 1901.

Collected by: R.S. Pieter Faure.

#### Family Myochamidae

*Myodora rectangulata* Barnard, 1964b: 28

= *Myodora (Myodora) quadrata* E. A. Smith: Kilburn 1973: 709.

Syntypes: SAM-A9556. Twelve valves.

Locality: off O'Neil Peak, Natal; 92 m; 28 February 1901.

Collected by: R.S. Pieter Faure.

#### REFERENCES

- ALLANSON, B. 1958. On the systematics and distribution of the molluscan genus *Siphonaria* in South Africa. *Hydrobiologia* 12: 149-180.
- ASHBY, E. 1928. South African chitons, being a description of the Polyplacophora represented in the Turton Collection. *Proc. malac. Soc. Lond.* 18: 76-92.
- ASHBY, E. 1931. Monograph of the South African Polyplacophora. *Ann. S. Afr. Mus.* 30: 1-59.
- BABA, K. 1955. *Opisthobranchia of Sagami Bay. Supplement*. Tokyo: Iwanami Shoten.

- BARNARD, K. H. 1927. South African nudibranch Mollusca, with descriptions of new species, and a note on some specimens from Tristan d'Acunha. *Ann. S. Afr. Mus.* **25**: 171-215.
- BARNARD, K. H. 1934. A new species of *Goniodoris* from South Africa. *J. Conch. Lond.* **20**: 2-3.
- BARNARD, K. H. 1957. The radula of '*Latiaxis fritschi*', and description of a new fossarid. *J. Conch. Lond.* **24**: 180-181.
- BARNARD, K. H. 1958. Contributions to the knowledge of South African marine Mollusca. Part I. Gastropoda: Prosobranchiata: Toxoglossa. *Ann. S. Afr. Mus.* **46**: 73-163.
- BARNARD, K. H. 1959. Contributions to the knowledge of South African marine Mollusca. Part II. Gastropoda: Prosobranchiata: Rhachiglossa. *Ann. S. Afr. Mus.* **45**: 1-237.
- BARNARD, K. H. 1960. New species of South African marine gastropods. *J. Conch. Lond.* **24**: 438-442.
- BARNARD, K. H. 1962a. A new genus in the family Marginellidae. *Proc. malac. Soc. Lond.* **35**: 14-15.
- BARNARD, K. H. 1962b. New species and records of South African marine Mollusca from Natal, Zululand and Mozambique. *Ann. Natal Mus.* **15**: 247-254.
- BARNARD, K. H. 1963a. Contributions to the knowledge of South African marine Mollusca. Part III. Gastropoda: Prosobranchiata: Taenioglossa. *Ann. S. Afr. Mus.* **47**: 1-199. (Published February.)
- BARNARD, K. H. 1963b. Deep-sea Mollusca from the region south of Madagascar. *Investl. Rep. Div. Sea Fish. Rep. S. Afr.* **44**: 3-19. (Published April.)
- BARNARD, K. H. 1963c. Deep-sea Mollusca from west of Cape Point, South Africa. *Ann. S. Afr. Mus.* **46**: 407-452. (Published April.)
- BARNARD, K. H. 1963d. Contributions to the knowledge of South African marine Mollusca. Part IV. Gastropoda: Prosobranchiata: Rhipidoglossa, Docoglossa. Tectibranchiata. Polyplacophora. Solenogastres. Scaphopoda. *Ann. S. Afr. Mus.* **47**: 201-360. (Published December.)
- BARNARD, K. H. 1964a. Contributions to the knowledge of South African marine Mollusca. Part V. Lamellibranchiata. *Ann. S. Afr. Mus.* **47**: 361-593.
- BARNARD, K. H. 1964b. Two genera of Erycinacea (Bivalvia) from South Africa. *Proc. malac. Soc. Lond.* **36**: 33-37.
- BARNARD, K. H. 1964c. The work of the S.S. *Pieter Faure* in Natal waters, with special reference to the Crustacea and Mollusca with descriptions of new species of Mollusca from Natal. *Ann. Natal Mus.* **16**: 9-29.
- BARNARD, K. H. 1969. Contributions to the knowledge of South African marine Mollusca. Part VI. Supplement. *Ann. S. Afr. Mus.* **47**: 595-661.
- BARNARD, K. H. 1974. Contributions to the knowledge of the South African marine Mollusca. Part VII. Revised fauna list. *Ann. S. Afr. Mus.* **47**: 663-781.
- BOSHOFF, P. 1968. Descriptions of three new species of Pelecypoda dredged off the coast of Natal, South Africa. *Trans. R. Soc. S. Afr.* **38**: 95-98.
- BOSS, K. 1969. The subfamily Tellininae in South African waters (Bivalvia, Mollusca). *Bull. Mus. comp. Zool. Harv.* **138**: 81-162.
- CATE, J. 1964. A new species of *Mitra* from the western Indian Ocean. *Veliger* **6**: 219-220.
- CERNOHORSKY, W. 1976. The Mitridae of the world. Part I. The subfamily Mitrinae. In: ABBOTT, R. T. ed. *Indo-Pacific Mollusca*: 273-528.
- GOSLINER, T. 1981. The South African Janolidae (Mollusca, Nudibranchia) with the description of a new genus and two new species. *Ann. S. Afr. Mus.* **86**: 1-42.
- GOSLINER, T. & GRIFFITHS, R. 1981. Description and revision of some South African aeolidacean Nudibranchia (Mollusca, Gastropoda) *Ann. S. Afr. Mus.* **84**: 105-150.
- GOSLINER, T. & LILTVED, W. 1982. Comparative morphology of three South African Triviidae (Gastropoda: Prosobranchia) with the description of a new species. *Zool. J. Linn. Soc.* **74**: 111-132.
- HOUBRICK, R. 1979. Classification and systematic relationships of the Abysochyrsidae, a relict family of bathyal snails (Prosobranchia: Gastropoda). *Smithson. Contr. Zool.* **290**: 1-21.
- KILBURN, R. 1971. On some species of the families Tonnidae, Hippocinidae, Buccinidae, Columbariidae, Fasciolariidae, Psammobiidae and Mactridae (Mollusca) in South African waters. *Ann. Natal Mus.* **20**: 483-497.

- KILBURN, R. 1973a. Notes on some Mollusca from Natal and Moçambique, with description of new species and subspecies of *Calliostoma*, *Solariella*, *Latiaxis*, *Babylonia*, *Fusinus*, *Bathytoma* and *Conus*. *Ann. Natal Mus.* **21**: 557–578.
- KILBURN, R. 1973b. The type of material of African marine Mollusca in the Natal Museum collection. Part I. Bivalvia. *Ann. Natal Mus.* **21**: 697–711.
- KILBURN, R. 1974. Taxonomic notes on South African marine Mollusca. Part 3. Gastropoda: Prosobranchia, with descriptions of new taxa of Naticidae, Fasciolaridae, Magilidae, Volutomitridae and Turridae. *Ann. Natal Mus.* **22**: 187–220.
- KILBURN, R. 1977. Taxonomic studies on the marine Mollusca of southern Africa and Mozambique. Part I. *Ann. Natal Mus.* **23**: 173–214.
- KILBURN, R. 1978. The Emarginulinae (Mollusca: Gastropoda: Fissurellidae) of southern Africa and Mozambique. *Ann. Natal Mus.* **23**: 431–453.
- KILBURN, R. 1981. Revision of the genus *Ancilla* Lamarck, 1799 (Mollusca: Olividae: Ancillinae). *Ann. Natal Mus.* **24**: 349–463.
- MACNAE, W. 1954. On some eolidacean nudibranchiate molluscs from South Africa. *Ann. Natal Mus.* **13**: 1–50.
- KNUDSEN, J. 1970. The systematics and biology of abyssal and hadal Bivalvia. *Galathea Rep.* **11**: 7–236.
- PRUVOT-FOL, A. 1934. Les opisthobranches de Quoy et Gaimard. *Archs Mus. Hist. Nat. Paris* (6) **11**: 13.
- RADWIN, G. & D'ATTILIO, A. 1976. *Murex shells of the world: an illustrated guide to the Muricidae*. Stanford: Stanford University Press.
- REHDER, H. 1969. *Voluticorbis* and *Fusivoluta*, two genera of deepwater Volutidae from South Africa. *Veliger* **11**: 200–209.
- SHACKELFORD, L. 1914. Two new species of *Marginella* from South Africa. *Ann. S. Afr. Mus.* **13**: 91–98.
- SHACKELFORD, L. 1916. Two new species of *Marginella* from South Africa. *Ann. S. Afr. Mus.* **13**: 193–194.
- SMITH, E. A. 1899. Descriptions of South African marine shells. *J. Conch. Lond.* **9**: 247–252.
- SMITH, E. A. 1901. On South African marine shells with descriptions of new species. *J. Conch. Lond.* **10**: 104–116.
- SOWERBY, G. B. 1894. Marine shells of South Africa. *J. Conch. Lond.* **7**: 368–378.
- SOWERBY, G. B. *Appendix to marine shells of South Africa*. London: Sowerby.
- SOWERBY, G. B. 1900. On some marine shells from Pondoland and the Kowie, with descriptions of seventeen new species. *Proc. malac. Soc. Lond.* **4**: 1–7.
- SOWERBY, G. B. 1901. On seven new species of marine Mollusca collected by Dr H. Becker at 'The Kowie', South Africa. *Proc. malac. Soc. Lond.* **4**: 213–215.
- SOWERBY, G. B. 1902. Mollusca of South Africa. *Mar. Invest. S. Afr.* **2**: 93–100.
- SOWERBY, G. B. 1903. Mollusca of South Africa. *Mar. Invest. S. Afr.* **2**: 213–232.
- SOWERBY, G. B. 1904. Mollusca of South Africa (Pelecypoda). *Mar. Invest. S. Afr.* **4**: 1–19.
- SOWERBY, G. B. 1906. On new species of *Siphonaria*, *Terebra* and *Mangilia*, and a remarkable form of *Cypraea cruenta*, from South Africa. *Proc. malac. Soc. Lond.* **7**: 37–39.
- SOWERBY, G. B. 1916. Descriptions of two new Mollusca of the genera *Leptothyra* and *Mitra*. *Ann. Mag. nat. Hist.* (8) **18**: 491–492.
- TARDY, J. 1969. Un nouveau genre de nudibranche méconnu des côtes Atlantique et de la Manche: *Pruvotfolia* (nov. g.) *pselliotes* (Labbé, 1923). *Vie Milieu* **20**: 327–346.
- THIELE, J. 1925. Gastropoda der Deutschen Tiefsee—Expedition. Teil 2. *Wiss. Ergebn. 'Valdiva'* **17**: 37–382.
- TOMLIN, J. R. 1920. Four new marine species from South Africa. *J. Conch. Lond.* **16**: 87–88.
- TOMLIN, J. R. 1921. Six new marine shells from South Africa. *J. Conch. Lond.* **16**: 215–217.
- TOMLIN, J. R. 1925. Reports on the marine Mollusca in the collections of the South African Museum. I. Turritellidae. *Ann. S. Afr. Mus.* **20**: 309–316.
- TOMLIN, J. R. 1927. Reports on the marine Mollusca in the collections of the South African Museum. II. Families Abysochrysidae, Oocorythidae, Haliotidae, Tonnidae. *Ann. S. Afr. Mus.* **25**: 77–83.
- TOMLIN, J. R. 1928. Reports on the marine Mollusca in the collections of the South African Museum III. Revision of the South African Nassariidae (*olim* Nassidae). IV. Families Terebridae, Columbariidae, Thaididae, Architectonicidae. *Ann. S. Afr. Mus.* **25**: 313–335.

- TOMLIN, J. R. 1931. Reports on the marine Mollusca in the collections of the South African Museum. V. The Scaphopoda. *Ann. S. Afr. Mus.* **29**: 337-340.
- TOMLIN, J. R. 1932. Reports on the marine Mollusca in the collections of the South African Museum. VI-VIII. *Ann. S. Afr. Mus.* **30**: 157-169.
- TOMLIN, J. R. 1939. A new South African vermetid. *J. Conch. Lond.* **21**: 145.
- TOMLIN, J. R. 1944. New South African *Siphonaria*. *J. Conch. Lond.* **22**: 92-93.
- TOMLIN, J. R. 1948. A new species of *Pleurotomaria*. *J. Conch. Lond.* **23**: 2.
- WALLS, J. 1979. *Cone shells, a synopsis of the living Conidae*. Hong Kong: T. F. H. Publications.
- WEAVER, C. 1963. Volute problems. *Hawaii. Shell News* **11** (5): 5.
- WEAVER, C. & DUPONT, J. 1970. Living volutes. A monograph of the recent Volutidae of the world. *Delaware Museum of Natural History. Monographic Ser.* **1**: 1-375.